

A generalization of twisted modules over vertex algebras

Kenichiro Tanabe*

Department of Mathematics
Hokkaido University
Kita 10, Nishi 8, Kita-Ku, Sapporo, Hokkaido, 060-0810
Japan

ktanabe@math.sci.hokudai.ac.jp

Abstract

We introduce a notion of a (V, T) -module over a vertex algebra V for an arbitrary positive integer T , which is a generalization of a twisted V -module. Under some conditions on V , we construct an associative algebra $A_m^T(V)$ for $m \in (1/T)\mathbb{N}$ and an $A_m^T(V)$ - $A_n^T(V)$ -bimodule $A_{n,m}^T(V)$ for $n, m \in (1/T)\mathbb{N}$ and we establish a one-to-one correspondence between the set of isomorphism classes of simple left $A_0^T(V)$ -modules and that of simple $(1/T)\mathbb{N}$ -graded (V, T) -modules.

1 Introduction

Twisted modules (or twisted sectors) were introduced in the study of the so-called orbifold models of conformal field theory (cf.[2, 3]). Let V be a vertex operator algebra and G a finite automorphism group of V . In terms of vertex operator algebras, the study of the orbifold models corresponds to the study of the subalgebra V^G of G -invariants in V .

*Partially supported by JSPS Grant-in-Aid for Scientific Research No. 20740002.

One of the main problems about V^G is to describe the V^G -modules in terms of V and G . Twisted modules have been studied systematically as representations of V related to this problem (cf. [6, 10, 12, 14]). For $g \in G$, every g -twisted V -module becomes a V^G -module. Moreover, it is conjectured that under some conditions on V , every simple V^G -module is contained in some simple g -twisted V -module for some $g \in G$ (cf.[3]). However, the following easy observation tells us an inconvenience of twisted V -modules from the representation theoretic viewpoint: let g, h be two different elements of G , M a g -twisted V -module and N an h -twisted V -module. Although $M \oplus N$ is a V^G -module, this is not a (twisted) V -module in general. This is one of obstructions to develop the representation theory of V^G .

In this paper, we first introduce a notion of a (V, T) -module for a vertex algebra V and a positive integer T , which is a generalization of a twisted V -module, in order to resolve the inconvenience just mentioned above. We next generalize the results by Zhu[16] to (V, T) -modules. In [16], if V is a vertex operator algebra, then Zhu constructed an associative algebra $A(V)$ and gave a one-to-one correspondence between the set of isomorphism classes of the simple $A(V)$ -modules and that of the simple V -modules with some conditions. Some generalizations of $A(V)$ have been obtained in [4, 5, 6, 7, 8] and they have played an important role in the representation theory of V . We shall show the following results for a vertex algebra V with a grading $V = \bigoplus_{i=\Delta}^{\infty} V_i$ such that $\Delta \in \mathbb{Z}_{\leq 0}$, $\mathbf{1} \in V_0$ and for all homogeneous element $a \in V$, $a_i V_j \subset V_{\text{wt } a - 1 - i + j}$, where $V_i = 0$ for $i < \Delta$. For every positive integer T and $n, m \in (1/T)\mathbb{N}$, we shall construct an associative algebra $A_m^T(V)$ and an $A_n^T(V)$ - $A_m^T(V)$ -bimodule $A_{n,m}^T(V)$ in Theorem 4.5. If $T = 1$, then $A_{n,m}^T(V)$ is the same as $A_{n,m}(V)$ in [5] and $A_n^T(V)$ is the same as $A_n(V)$ in [7]. In particular, $A_0^1(V)$ is the same as $A(V)$ in [16]. For an automorphism g of V of finite order, $A_{g,n,m}(V)$ in [6, 8] is a quotient of $A_{n,m}^{|g|}(V)$. For $m \in (1/T)\mathbb{N}$ and a left $A_m^T(V)$ -module U , we shall show in Theorem 5.13 that the $(1/T)\mathbb{N}$ -graded vector space $M(U) = \bigoplus_{n \in (1/T)\mathbb{N}} A_{n,m}^T(V) \otimes_{A_m^T(V)} U$ has a structure of (V, T) -module with a universal property. In Corollary 5.14, we establish a one-to-one correspondence between the set of isomorphism classes of simple $A_0^T(V)$ -modules and that of simple $(1/T)\mathbb{N}$ -graded (V, T) -modules.

The organization of the paper is as follows. In Section 2 we introduce a notion of a (V, T) -module. In Section 3 we introduce a subspace $O_{n,m}^{T,1}(\alpha, \beta; x)$ of $\mathbb{C}[z, z^{-1}]$ for $n, m \in (1/T)\mathbb{N}$ and $\alpha, \beta \in \mathbb{Z}$ and

study its properties.. In Section 4 we construct an associative algebra $A_m^T(V)$ and an $A_n^T(V)$ - $A_m^T(V)$ -bimodule $A_m^T(V)$ for $n, m \in (1/T)\mathbb{N}$ by using the results in Section 4. In Section 5 we introduce a notion of a $(1/T)\mathbb{N}$ -graded (V, T) -module and study a relation between the $(1/T)\mathbb{N}$ -graded (V, T) -modules and the left $A_m^T(V)$ -modules. Section 6 consists of two subsections. In Subsection 6.1 we compute the determinant of a matrix used in Section 3. In Subsection 6.2 we improve some results in [15].

2 (V, T) -modules

We assume that the reader is familiar with the basic knowledge on vertex algebras as presented in [1, 6, 13].

Throughout this paper, $\mathbb{N}$ denotes the set of all non-negative integers, T is a fixed positive integer and $(V, Y, \mathbf{1})$ is a vertex algebra. Recall that V is the underlying vector space, $Y(-, x)$ is the linear map from V to $(\text{End } V)[[x, x^{-1}]]$, and $\mathbf{1}$ is the vacuum vector. For $i, j \in \mathbb{Z}$, define

$$\begin{aligned}\mathbb{Z}_{\leq i} &= \{k \in \mathbb{Z} \mid k \leq i\}, \\ \mathbb{Z}_{\geq i} &= \{k \in \mathbb{Z} \mid k \geq i\}, \\ \mathbb{C}[x, x^{-1}]_{\leq i} &= \text{Span}_{\mathbb{C}}\{x^k \mid k \leq i\}, \\ \mathbb{C}[x, x^{-1}]_{j, i} &= \text{Span}_{\mathbb{C}}\{x^k \mid j \leq k \leq i\}.\end{aligned}$$

For $f(z) \in \mathbb{C}[z, z^{-1}]$ and $a, b \in V$, $f(z)|_{z^j=a_jb}$ denotes the element of V obtained from $f(z)$ by replacing z^j by a_jb for all $j \in \mathbb{Z}$. For $i, j \in \mathbb{Q}$, define

$$\delta(i \leq j) = \begin{cases} 1 & \text{if } i \leq j, \\ 0 & \text{if } i > j. \end{cases} \quad (2.1)$$

Let M be a vector space over $\mathbb{C}$. Define three linear injective maps

$$\begin{aligned}\iota_{x,y} : M[[x^{1/T}, y^{1/T}]] [x^{-1/T}, y^{-1/T}, (x-y)^{-1}] &\rightarrow M((x^{1/T}))((y^{1/T})), \\ \iota_{y,x} : M[[x^{1/T}, y^{1/T}]] [x^{-1/T}, y^{-1/T}, (x-y)^{-1}] &\rightarrow M((y^{1/T}))((x^{1/T})), \\ \iota_{x,x-y} : M[[x^{1/T}, y^{1/T}]] [x^{-1/T}, y^{-1/T}, (x-y)^{-1}] &\rightarrow M((y^{1/T}))((x-y))\end{aligned}$$

by

$$\begin{aligned}\iota_{x,y}f &= \sum_{j,k,l} a_{j,k,l} \sum_{i=0}^{\infty} \binom{l}{i} (-1)^i x^{j+l-i} y^{k+i}, \\ \iota_{y,x}f &= \sum_{j,k,l} a_{j,k,l} \sum_{i=0}^{\infty} \binom{l}{i} (-1)^{l-i} y^{k+l-i} x^{j+i}, \\ \iota_{y,x-y}f &= \sum_{j,k,l} a_{j,k,l} \sum_{i=0}^{\infty} \binom{j}{i} y^{k+j-i} (x-y)^{l+i}\end{aligned}$$

for $f = \sum_{j,k,l} a_{j,k,l} x^j y^k (x-y)^l \in M[[x^{1/T}, y^{1/T}]] [x^{-1/T}, y^{-1/T}, (x-y)^{-1}]$, $a_{i,j,k} \in M$. We can also define the map

$$\iota_{x-y,y} : M[[x^{1/T}, y^{1/T}]] [x^{-1/T}, y^{-1/T}, (x-y)^{-1}] \rightarrow M(((x-y)^{1/T}))((y^{1/T}))$$

similarly. Since $\iota_{x,y}(x-y)^i = \sum_{j=0}^{\infty} \binom{i}{j} x^{i-j} (-1)^j y^j$ and $\iota_{x-y,y} x^i = \sum_{j=0}^{\infty} \binom{i}{j} (x-y)^{i-j} y^j$, we identify $M(((x-y)^{1/T}))((y^{1/T}))$ with $M((x^{1/T}))((y^{1/T}))$ and $\iota_{x-y,y}$ with $\iota_{x,y}$.

Now we introduce a generalization of a twisted module.

Definition 2.1. Let M be a vector space over $\mathbb{C}$ and $Y_M(-, x)$ be a linear map from V to $(\text{End}_{\mathbb{C}} M)[[x^{1/T}, x^{-1/T}]]$. We call (M, Y_M) a (V, T) -module if

- (1) For $a \in V$ and $w \in M$, $Y_M(a, x)w \in M((x^{1/T}))$.
- (2) $Y_M(\mathbf{1}, x) = \text{id}_M$.
- (3) For $a, b \in V$ and $w \in M$, there is $F(a, b, w|x, y) \in M[[x^{1/T}, y^{1/T}]] [x^{-1/T}, y^{-1/T}, (x-y)^{-1}]$ such that

$$\begin{aligned}\iota_{x,y}F(a, b, w|x, y) &= Y_M(a, x)Y_M(b, y)w, \\ \iota_{y,x}F(a, b, w|x, y) &= Y_M(b, y)Y_M(a, x)w, \quad \text{and} \\ \iota_{y,x-y}F(a, b, w|x, y) &= Y_M(Y(a, x-y)b, y)w.\end{aligned}$$

We note that in Definition 2.1, $F(a, b, w|x, y)$ is uniquely determined by $a, b \in V$ and $w \in M$ since $\iota_{x,y}$ is an injection. For a (V, T) -module M , a subspace N of M is called (V, T) -submodule of M if $(N, Y_M|_N)$ is a (V, T) -module, where $Y_M|_N$ is the restriction of Y_M to N . A non-zero (V, T) -module M is called *simple* if there is no submodule of M except 0 and M itself. For a submodule N of a (V, T) -module M , the quotient space M/N is clearly a (V, T) -module. For a set of (V, T) -modules $\{M_i\}_{i \in I}$, the direct sum $\oplus_{i \in I} M_i$ is a (V, T) -module.

Remark 2.2. It follows from Lemma 2.4 below that every $(V, 1)$ -module is a V -module and vice versa and that every g -twisted V -module is a $(V, |g|)$ -module for an automorphism g of V of finite order. Let T' be a positive multiple of T . Then every (V, T) -module is a (V, T') -module.

Example 2.3. We introduce an easy example of simple (V, T) -modules which is not a twisted V -module. Let V be a simple vertex operator algebra. Suppose the symmetric group S_3 of degree 3 is an automorphism group of V . Let $\sigma, \tau \in S_3$ such that $|\sigma| = 3$ and $|\tau| = 2$ and $M = \bigoplus_{j \in (1/3)\mathbb{N}} M(j)$ a simple σ -twisted V -module [6]. It follows from Remark 2.2 that M is a $(V, 3)$ -module. Restricting Y_M to $V^{(\tau)}$, M becomes a $(V^{(\tau)}, 3)$ -module. We shall show M is a simple $(V^{(\tau)}, 3)$ -module. Let W be a non-zero $(V^{(\tau)}, 3)$ -submodule of M . We denote the subspace $\bigoplus_{j \in i/3 + \mathbb{N}} M(j)$ of M by $M^i, i = 0, 1, 2$. Since $\tau\sigma\tau = \sigma^{-1} \neq \sigma$, an improvement of [15, Theorem 2] (see Subsection 6.2) implies that M^0, M^1 and M^2 are all inequivalent simple V^{S_3} -modules. Thus, W contains at least one of M^0, M^1 and M^2 since $V^{S_3} \subset V^{(\tau)}$. We denote the eigenspace $\{v \in V \mid \sigma v = e^{-2\pi\sqrt{-1}r/3}v\}$ of σ by $V^{(\sigma, r)}, r = 0, 1, 2$. It follows by [9, Proposition 3.3] and [11, Theorem 1] that $V^{(\tau)} \not\subset V^{(\sigma)}$ and hence there exists $a = a^0 + a^1 + a^2 \in V^{(\tau)}$, $a^r \in V^{(\sigma, r)}$ such that at least one of a^1, a^2 is not zero. Since

$$Y_M(a, x) = \sum_{i \in \mathbb{Z}} a_i^0 x^{-i-1} + \sum_{i \in 1/3 + \mathbb{Z}} a_i^1 x^{-i-1} + \sum_{i \in 2/3 + \mathbb{Z}} a_i^2 x^{-i-1}$$

and M is a simple σ -twisted V -module, W contains at least two of M^0, M^1 and M^2 . Repeating the same argument, we obtain that M is a simple $(V^{(\tau)}, 3)$ -module.

Since at least one of a^1, a^2 above is not zero, M is not a V -module. Suppose M is a g -twisted $V^{(\tau)}$ -module for some $g \in \text{Aut } V^{(\tau)}$ of order 3. Then, the eigenspace $(V^{(\tau)})^{(g, r)} = \{v \in V^{(\tau)} \mid gv = e^{-2\pi\sqrt{-1}r/3}v\}$ of g is a subspace of $V^{(\sigma, r)}$ for each $r = 0, 1, 2$ since $Y_M(b, x) = \sum_{j \in r/3 + \mathbb{Z}} b_j x^{-j-1}$ for $b \in (V^{(\tau)})^{(g, r)}$. Therefore, $(V^{(\tau)})^{(g, 1)} = (V^{(\tau)})^{(g, 2)} = 0$ since there is no representation ρ of S^3 such that $\rho(\sigma) = e^{-2\pi\sqrt{-1}r/3}$ and $\rho(\tau) = 1$ for $r = 1, 2$. This contradicts to that the order of g is equal to 3. We conclude that M is not a twisted V -module.

Let M be a vector space. For $r = 0, \dots, T-1$ and $X(x, y) = \sum_{i, j \in (1/T)\mathbb{Z}} X_{ij} x^i y^j \in M[[x^{1/T}, x^{-1/T}, y^{1/T}, y^{-1/T}]]$, $X_{ij} \in M$, we de-

fine

$$\begin{aligned}
X(x, y)^{r, x} &= \sum_{\substack{i \in r/T + \mathbb{Z} \\ j \in (1/T)\mathbb{Z}}} X_{ij} x^i y^j \quad \text{and} \\
X(x, y)^{r, y} &= \sum_{\substack{i \in (1/T)\mathbb{Z} \\ j \in r/T + \mathbb{Z}}} X_{ij} x^i y^j.
\end{aligned} \tag{2.2}$$

In the same way, we define $X(x, y)^{r, x}$ and $X(x, y)^{r, y}$ for $X(x, y) \in M[[x_1^{1/T}, x_2^{1/T}]] [x_1^{-1/T}, x_2^{-1/T}, (x_1 - x_2)^{-1}]$. Clearly, $\sum_{r=0}^{T-1} X(x, y)^{r, x} = \sum_{r=0}^{T-1} X(x, y)^{r, y} = X(x, y)$. For $0 \leq s \leq T-1$, $j \in s/T + \mathbb{Z}$, $k \in (1/T)\mathbb{Z}$ and $l \in (1/T)\mathbb{Z}$, the following fact is well known and straightforward:

$$\begin{aligned}
& x_0^{-1} \delta\left(\frac{x_1 - x_2}{x_0}\right) \iota_{x_1, x_2}((x_1^j x_2^k x_0^l)|_{x_0=x_1-x_2}) \\
& - x_0^{-1} \delta\left(\frac{-x_2 + x_1}{x_0}\right) \iota_{x_2, x_1}((x_1^j x_2^k x_0^l)|_{x_0=x_1-x_2}) \\
& = x_1^{-1} \left(\frac{x_2 + x_0}{x_1}\right)^{-s/T} \delta\left(\frac{x_2 + x_0}{x_1}\right) \iota_{x_2, x_0}((x_1^j x_2^k x_0^l)|_{x_1=x_2+x_0}).
\end{aligned} \tag{2.3}$$

The argument in the proof of the following lemma is well known (cf. [13, Sections 3.2–3.4]).

Lemma 2.4. *Let $A(x_1, x_2) \in M((x_1^{1/T}))((x_2^{1/T}))$, $B(x_2, x_1) \in M((x_2^{1/T}))((x_1^{1/T}))$, and $C(x_2, x_0) \in M((x_2^{1/T}))((x_0))$. Then, the three following conditions are equivalent.*

(1) *There is $F \in M[[x_1^{1/T}, x_2^{1/T}]] [x_1^{-1/T}, x_2^{-1/T}, (x_1 - x_2)^{-1}]$ such that*

$$\begin{aligned}
\iota_{x_1, x_2} F &= A(x_1, x_2), & \iota_{x_2, x_1} F &= B(x_2, x_1), \quad \text{and} \\
\iota_{x_2, x_1-x_2} F &= C(x_2, x_1 - x_2).
\end{aligned}$$

(2) *There are $C^{[s]}(x_2, x_0) \in M((x_2^{1/T}))((x_0))$, $s = 0, \dots, T-1$ such that $\sum_{s=0}^{T-1} C^{[s]}(x_2, x_0) = C(x_2, x_0)$ and*

$$\begin{aligned}
& x_0^{-1} \delta\left(\frac{x_1 - x_2}{x_0}\right) A(x_1, x_2)^{s, x_1} - x_0^{-1} \delta\left(\frac{-x_2 + x_1}{x_0}\right) B(x_2, x_1)^{s, x_1} \\
& = x_1^{-1} \left(\frac{x_2 + x_0}{x_1}\right)^{-s/T} \delta\left(\frac{x_2 + x_0}{x_1}\right) C^{[s]}(x_2, x_0).
\end{aligned} \tag{2.4}$$

(3) There are positive integers l, q and $C^{[s]}(x_2, x_0) \in M((x_2^{1/T}))((x_0))$, $s = 0, \dots, T-1$ such that $\sum_{s=0}^{T-1} C^{[s]}(x_2, x_0) = C(x_2, x_0)$,

$$(x_1 - x_2)^l A(x_1, x_2) = (x_1 - x_2)^l B(x_2, x_1) \quad (2.5)$$

in $M[[x_1^{1/T}, x_2^{1/T}, x_1^{-1/T}, x_2^{-1/T}]]$ and

$$\begin{aligned} & \iota_{x_0, x_2}(x_0 + x_2)^{-s/T+q} (A(x_1, x_2)^{s, x_1})|_{x_1=x_0+x_2} \\ &= \iota_{x_2, x_0}(x_2 + x_0)^{-s/T+q} C^{[s]}(x_2, x_0) \end{aligned} \quad (2.6)$$

in $M[[x_0, x_2^{1/T}, x_0^{-1}, x_2^{-1/T}]]$.

In this case, F and $C^{[s]}(x_2, x_0)$, $s = 0, \dots, T-1$ are uniquely determined by $A(x_1, x_2)$, $B(x_2, x_1)$ and $C(x_2, x_0)$.

Proof. We show (1) implies (2). Define $C^{[s]}(x_2, x_0) \in M((x_2^{1/T}))((x_0))$ by $C^{[s]}(x_2, x_1 - x_2) = \iota_{x_2, x_1-x_2} F^{s, x_1} \in M((x_2^{1/T}))((x_1 - x_2))$ for $s = 0, \dots, T-1$. Clearly, $\sum_{s=0}^{T-1} C^{[s]}(x_2, x_0) = C(x_2, x_0)$. Since $\iota_{x_1, x_2} F^{s, x_1} = A(x_1, x_2)^{s, x_1}$ and $\iota_{x_2, x_1} F^{s, x_1} = B(x_2, x_1)^{s, x_1}$ for $s = 0, \dots, T-1$, (2.4) follows from (2.3).

We show (2) implies (3). Let l be a positive integer such that $x_0^l C^{[s]}(x_2, x_0) \in M((x_2^{1/T}))[[x_0]]$ for all $s = 0, \dots, T-1$. Multiplying (2.4) by x_0^l and then taking Res_{x_0} , we have $(x_1 - x_2)^l A(x_1, x_2)^{s, x_1} = (x_1 - x_2)^l B(x_2, x_1)^{s, x_1}$ and hence (2.5). Let q be a positive integer such that $x_1^{-s/T+q} B(x_2, x_1)^{s, x_1} \in M((x_2^{1/T}))[[x_1]]$ for all $s = 0, \dots, T-1$. Multiplying (2.4) by $x_1^{-s/T+q}$ and then taking Res_{x_1} , we have (2.6).

We show (3) implies (1). Since the left-hand side of (2.5) is an element of $M((x_1^{1/T}))((x_2^{1/T}))$ and the right-hand side of (2.5) is an element of $M((x_2^{1/T}))((x_1^{1/T}))$, $G = (x_1 - x_2)^l A(x_1, x_2) = (x_1 - x_2)^l B(x_2, x_1)$ is an element of $M[[x_1^{1/T}, x_2^{1/T}]] [x_1^{-1/T}, x_2^{-1/T}]$. Define

$$F = (x_1 - x_2)^{-l} G \in M[[x_1^{1/T}, x_2^{1/T}]] [x_1^{-1/T}, x_2^{-1/T}, (x_1 - x_2)^{-1}].$$

It is clear that $\iota_{x_1, x_2} F = A(x_1, x_2)$ and $\iota_{x_2, x_1} F = B(x_2, x_1)$. Applying the same argument to (2.6), we obtain $H_s \in M[[x_1^{1/T}, x_2^{1/T}]] [x_1^{-1/T}, x_2^{-1/T}, (x_1 - x_2)^{-1}]$, $s = 0, \dots, T-1$ such that

$$\begin{aligned} \iota_{x_1-x_2, x_2} H_s &= A(x_1, x_2)^{s, x_1} = A((x_1 - x_2) + x_2, x_2)^{s, x_1} \text{ and} \\ \iota_{x_2, x_1-x_2} H_s &= C^{[s]}(x_1, x_1 - x_2). \end{aligned}$$

Since $M(((x_1 - x_2)^{1/T}))((x_2^{1/T})) = M((x_1^{1/T}))((x_2^{1/T}))$ and ι_{x_1, x_2} is injective, we have $F^{s, x_1} = H_s$ for all $s = 0, \dots, T-1$ and therefore $\iota_{x_2, x_1-x_2} F = C(x_2, x_1 - x_2)$.

We show F and $C^{[s]}(x_2, x_0)$, $s = 0, \dots, T-1$ are uniquely determined. Since ι_{x_1, x_2} is injective and $\iota_{x_1, x_2} F = A(x_1, x_2)$, F is uniquely determined. In the above argument that (3) implies (1), we have constructed F such that $\iota_{x_1, x_2} F = A(x_1, x_2)$ and $\iota_{x_2, x_1-x_2} F^{s, x_1} = C^{[s]}(x_2, x_1 - x_2)$. Thus, $C^{[s]}(x_2, x_0)$, $s = 0, \dots, T-1$ in (3) are uniquely determined. A similar argument shows that $C^{[s]}(x_2, x_0)$, $s = 0, \dots, T-1$ in (2) are uniquely determined and that $C^{[s]}(x_2, x_0)$ in (2) is the same as that in (3) for each s . $\square$

Remark 2.5. The following facts for (2.4) are well known and straightforward.

- (1) A direct computation shows that (2.4) is equivalent to

$$\begin{aligned} & x_0^{-1} \delta\left(\frac{x_1 - x_2}{x_0}\right) A(x_1, x_2) - x_0^{-1} \delta\left(\frac{-x_2 + x_1}{x_0}\right) B(x_2, x_1) \\ &= \frac{1}{T} \sum_{j=0}^{T-1} x_1^{-1} \delta(e^{2\pi\sqrt{-1}j/T} \frac{(x_2 + x_0)^{1/T}}{x_1^{1/T}}) \sum_{s=0}^{T-1} e^{2\pi\sqrt{-1}js/T} C^{[s]}(x_2, x_0). \end{aligned} \quad (2.7)$$

- (2) If we write $A(x_1, x_2) = \sum_{p,q} A_{(p,q)} x_1^{-p-1} x_2^{-q-1}$, $B(x_2, x_1) = \sum_{p,q} B_{(p,q)} x_2^{-p-1} x_1^{-q-1}$ and $C^{[s]}(x_2, x_0) = \sum_{p,q} C_{(p,q)}^{[s]} x_2^{-p-1} x_0^{-q-1}$, where $A_{(p,q)}, B_{(p,q)}, C_{(p,q)} \in M$, then we have

$$\begin{aligned} & \sum_{i=0}^{\infty} \binom{l}{i} (-1)^i (A_{(l+j-i, k+i)} + (-1)^{l+1} B_{(l+k-i, j+i)}) \\ &= \sum_{i=0}^{\infty} \binom{j}{i} C_{(j+k-i, l+i)}^{[-s]} \end{aligned} \quad (2.8)$$

for $0 \leq s \leq T-1$, $j \in s/T + \mathbb{Z}$, $k \in (1/T)\mathbb{Z}$ and $l \in \mathbb{Z}$ by comparing the coefficients of both sides of (2.4). Thus, a direct computation shows that (2.4) is also equivalent to the condition that

$$\begin{aligned} & \text{Res}_{x_1} A(x_1, x_2) \iota_{x_1, x_2} (x_1^j x_2^k (x_1 - x_2)^l) \\ & \quad - \text{Res}_{x_1} B(x_2, x_1) \iota_{x_2, x_1} (x_1^j x_2^k (x_1 - x_2)^l) \\ &= \text{Res}_{x_1-x_2} C^{[-s]}(x_2, x_1 - x_2) \iota_{x_2, x_1-x_2} (x_1^j x_2^k (x_1 - x_2)^l) \end{aligned} \quad (2.9)$$

in $M[[x_2^{1/T}, x_2^{-1/T}]]$ for all $0 \leq s \leq T-1$, $j \in s/T + \mathbb{Z}$, $k \in (1/T)\mathbb{Z}$ and $l \in \mathbb{Z}$. Here, Res_x is defined by

$$\text{Res}_x f(x) = f_{-1}$$

for $f(x) = \sum_{i \in (1/T)\mathbb{Z}} f_i x^i \in M[[x^{1/T}, x^{-1/T}]]$.

Remark 2.6. For $q \in \mathbb{Z}$ we denote by $M((x_2^{1/T}))((x_0))_{\geq q}$ the set of all elements in $M((x_2^{1/T}))((x_0))$ of the form $\sum_{\substack{i \in (1/T)\mathbb{Z} \\ j \in \mathbb{Z}_{\geq q}}} X_{ij} x_2^i x_0^j$. Suppose

$C(x_2, x_0)$ in Lemma 2.4 is an element of $M((x_2^{1/T}))((x_0))_{\geq q}$. Since $\iota_{x_2, x_1-x_2} x_1^j x_2^k (x_1 - x_2)^l = \sum_{i=0}^{\infty} \binom{j}{i} x_2^{k+j-i} (x_1 - x_2)^{l+i}$, we see that F in Lemma 2.4 (1) has the form $F = (x_1 - x_2)^q G$, where $G \in M[[x_1^{1/T}, x_2^{1/T}]][[x_1^{-1/T}, x_2^{-1/T}]]$. Thus, $C^{[s]}(x_2, x_0) = \iota_{x_2, x_1-x_2} F^{s, x_1} \in M((x_2^{1/T}))((x_0))_{\geq q}$ for all $s = 0, \dots, T-1$.

Let M be a (V, T) -module. For $a \in V$ and $s = 0, \dots, T-1$, we define $Y_M^s(a, x)$ by

$$Y_M^s(a, x) = \sum_{i \in s/T + \mathbb{Z}} a_i x^{-i-1}. \quad (2.10)$$

Let $a, b \in V$ and $w \in M$. We denote by $Y_M^{(-s)}(a, b|x_1, x_0)(w)$ the element $C^{[s]}(x_2, x_0)$ of $M((x_2^{1/T}))((x_0))$, $s = 0, \dots, T-1$ obtained by applying Lemma 2.4 to $A(x_1, x_2) = Y_M(a, x_1)Y_M(b, x_2)w$, $B(x_2, x_1) = Y_M(b, x_2)Y_M(a, x_1)w$ and $C(x_2, x_0) = Y_M(Y(a, x_0)b, x_2)w$. That is,

$$Y_M^{(s)}(a, b|x_1, x_1 - x_2)(w) = \iota_{x_2, x_1-x_2}(F(a, b, w|x_1, x_2)^{-s, x_1}). \quad (2.11)$$

The conditions in Lemma 2.4 (2) become

$$\sum_{s=0}^{T-1} Y_M^{(s)}(a, b|x_2, x_0)(w) = Y_M(Y(a, x_0)b, x_2)w \quad (2.12)$$

and

$$\begin{aligned} & x_0^{-1} \delta\left(\frac{x_1 - x_2}{x_0}\right) Y_M^s(a, x_1) Y_M(b, x_2) w - x_0^{-1} \delta\left(\frac{-x_2 + x_1}{x_0}\right) Y_M(b, x_2) Y_M^s(a, x_1) w \\ &= x_1^{-1} \left(\frac{x_2 + x_0}{x_1}\right)^{s/T} \delta\left(\frac{x_2 + x_0}{x_1}\right) Y_M^{(s)}(a, b|x_2, x_0)(w). \end{aligned} \quad (2.13)$$

The uniqueness of $F(a, b, w|x_1, x_2)$ for each $a, b \in V$ and $w \in M$ implies that for fixed $a, b \in V$ the map $Y_M^{(s)}(a, b|x_2, x_0) : M \rightarrow$

$M((x_2^{1/T}))((x_0))$ is linear and that the map $V \times V \ni (a, b) \mapsto Y_M^{(s)}(a, b|x_2, x_0) \in \text{Hom}_{\mathbb{C}}(M, M((x_2^{1/T}))((x_0)))$ is bilinear. We write

$$Y_M^{(s)}(a, b|x_2, x_0) = \sum_{i \in (1/T)\mathbb{Z}} \sum_{j \in \mathbb{Z}} Y_M^{(s)}(a, b; i, j) x_2^{-i-1} x_0^{-j-1},$$

where $Y_M^{(s)}(a, b; i, j) \in \text{End}_{\mathbb{C}} M$.

Remark 2.7. Let g be an automorphism of V of finite order, t a positive multiple of $|g|$ and (M, Y_M) a g -twisted V -module. As stated in Remark 2.2, (M, Y_M) is a (V, t) -module by Lemma 2.4. We explain what is $Y_M^{(s)}(a, b|x_2, x_0)$ for $a, b \in V$ and $s = 0, \dots, t-1$ in this case. We denote by $V^{(g,r)}$, $r = 0, \dots, t-1$ the eigenspace $\{v \in V \mid gv = e^{-2\pi\sqrt{-1}r/t}v\}$ of g . For $a \in V$, we denote by $a^{(g,r)}$ the r -th component of a in the decomposition $V = \bigoplus_{r=0}^{t-1} V^{(g,r)}$, that is, $a = \sum_{r=0}^{t-1} a^{(g,r)}$, $a^{(g,r)} \in V^{(g,r)}$.

Let $0 \leq s \leq t-1$, $a, b \in V$ and $w \in M$. Since

$$\begin{aligned} (Y_M(a, x_1)Y_M(b, x_2)w)^{-s, x_1} &= Y_M(a^{(g,s)}, x_1)Y_M(b, x_2)w \quad \text{and} \\ (Y_M(b, x_2)Y_M(a, x_1)w)^{-s, x_1} &= Y_M(b, x_2)Y_M(a^{(g,s)}, x_1)w, \end{aligned}$$

it follows by (2.4) that

$$Y_M^{(s)}(a, b|x_2, x_0)(w) = Y_M(Y(a^{(g,s)}, x_0)b, x_2)w. \quad (2.14)$$

Let $a, b \in V$, $w \in M$, $j, k \in (1/T)\mathbb{Z}$, $l \in \mathbb{Z}$ and s the integer uniquely determined by the conditions $0 \leq s \leq T-1$ and $s/T \equiv j \pmod{\mathbb{Z}}$. It follows by (2.8) or by comparing the coefficients of both sides of (2.13) that

$$\begin{aligned} &\sum_{i=0}^{\infty} \binom{j}{i} Y_M^{(s)}(a, b; j+k-i, l+i)(w) \\ &= \sum_{i=0}^{\infty} \binom{l}{i} (-1)^i (a_{l+j-i} b_{k+i} + (-1)^{l+1} b_{l+k-i} a_{j+i})w. \end{aligned} \quad (2.15)$$

It follows by (2.9) that

$$\begin{aligned} &\text{Res}_{x_1-x_2} \iota_{x_2, x_1-x_2} (x_1^j x_2^k (x_1-x_2)^l) Y_M^{(s)}(a, b|x_2, x_1-x_2)(w) \\ &= \text{Res}_{x_1} \iota_{x_1, x_2} (x_1^j x_2^k (x_1-x_2)^l) Y_M(a, x_1) Y_M(b, x_2)w \\ &\quad - \text{Res}_{x_1} \iota_{x_2, x_1} (x_1^j x_2^k (x_1-x_2)^l) Y_M(b, x_2) Y_M(a, x_1)w \\ &= \text{Res}_{x_1} \iota_{x_1, x_2} (x_1^j x_2^k (x_1-x_2)^l) Y_M^s(a, x_1) Y_M(b, x_2)w \\ &\quad - \text{Res}_{x_1} \iota_{x_2, x_1} (x_1^j x_2^k (x_1-x_2)^l) Y_M(b, x_2) Y_M^s(a, x_1)w. \end{aligned} \quad (2.16)$$

Lemma 2.8. *We use the notation above. Let L be an integer such that $a_i b = 0$ for all $i \in \mathbb{Z}_{\geq L+1}$. Then*

$$\begin{aligned} & Y_M^{(s)}(a, b; j+k, l)(w) \\ &= \sum_{m=0}^{L-l} \binom{-j}{m} \sum_{i=0}^{\infty} \binom{l+m}{i} (-1)^i (a_{l+m+j-i} b_{k-m+i} + (-1)^{l+m+1} b_{l+k-i} a_{j+i}) w \end{aligned} \quad (2.17)$$

Proof. It follows from Remark 2.6 that $Y_M^{(s)}(a, b|x_2, x_0)(w) \in M((x_2^{1/T}))((x_0))_{\geq -L-1}$. Thus, if $l > L$, then the both-sides of (2.17) are equal to 0. Suppose $l \leq L$. Define

$$R(m) = \sum_{i=0}^{\infty} \binom{m}{i} (-1)^i (a_{m+j-i} b_{k-m+l+i} + (-1)^{m+1} b_{l+k-i} a_{j+i}) w$$

for $m \in \mathbb{Z}_{\leq L}$. Since

$$\begin{aligned} & \begin{pmatrix} 1 & 0 & \cdots & \cdots & 0 \\ \binom{j}{1} & 1 & \ddots & & \vdots \\ \binom{j}{2} & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ \binom{j}{L-l} & \cdots & \binom{j}{2} & \binom{j}{1} & 1 \end{pmatrix} \begin{pmatrix} Y_M^{(s)}(a, b; j+k+l-L, L)(w) \\ Y_M^{(s)}(a, b; j+k+l-L+1, L-1)(w) \\ \vdots \\ Y_M^{(s)}(a, b; j+k, l)(w) \end{pmatrix} \\ &= \begin{pmatrix} R(L) \\ R(L-1) \\ \vdots \\ R(l) \end{pmatrix} \end{aligned}$$

by (2.15), we have

$$\begin{aligned} & \begin{pmatrix} Y_M^{(s)}(a, b; j+k+l-L, L)(w) \\ Y_M^{(s)}(a, b; j+k+l-L+1, L-1)(w) \\ \vdots \\ Y_M^{(s)}(a, b; j+k, l)(w) \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & \cdots & \cdots & 0 \\ \binom{-j}{1} & 1 & \ddots & & \vdots \\ \binom{-j}{2} & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ \binom{-j}{L-l} & \cdots & \binom{-j}{2} & \binom{-j}{1} & 1 \end{pmatrix} \begin{pmatrix} R(L) \\ R(L-1) \\ \vdots \\ R(l) \end{pmatrix}. \end{aligned}$$

This implies (2.17). $\square$

Let $a, b \in V$, $w \in M$, $j, k \in (1/T)\mathbb{Z}$, $l \in \mathbb{Z}$ and s the integer uniquely determined by the conditions $0 \leq s \leq T-1$ and $s/T \equiv j \pmod{\mathbb{Z}}$. It follows by Lemma 2.4 that $F(a, \mathbf{1}, w|x_1, x_2) = Y_M(a, x_1)$ since $Y_M(a, x_1)Y_M(\mathbf{1}, x_2)w = Y_M(a, x_1)w \in M[[x_1^{1/T}, x_2^{1/T}]]x_1^{-1/T}, x_2^{-1/T}, (x_1 - x_2)^{-1}]$. Comparing the coefficients of

$$\begin{aligned} & \iota_{x_2, x_1 - x_2} x_1^j Y_M^{(s)}(a, \mathbf{1}|x_2, x_1 - x_2)(w) \\ &= \sum_{k \in (1/T)\mathbb{Z}} \sum_{l \in \mathbb{Z}} \sum_{i=0}^{\infty} \binom{j}{i} Y_M^{(s)}(a, \mathbf{1}; j+k-i, l+i)(w) x_2^{-k-1} (x_1 - x_2)^{-l-1} \end{aligned}$$

and

$$\begin{aligned} & \iota_{x_2, x_1 - x_2} x_1^j Y_M^s(a, x_1)w \\ &= \sum_{k \in \mathbb{Z}} \sum_{l \in \mathbb{Z}} \binom{k}{-l-1} (-1)^{l+1} a_{j+k+l+1} w x_2^{-k-1} (x_1 - x_2)^{-l-1}, \end{aligned}$$

we have

$$\begin{aligned} & \sum_{i=0}^{-l-1} \binom{j}{i} Y_M^{(s)}(a, \mathbf{1}; j+k-i, l+i)(w) \\ &= \begin{cases} \binom{k}{-l-1} (-1)^{l+1} a_{j+k+l+1} w & \text{if } k \in \mathbb{Z}, \\ 0 & \text{if } k \notin \mathbb{Z}. \end{cases} \end{aligned} \quad (2.18)$$

Here, we used $Y_M^{(s)}(a, \mathbf{1}|x_2, x_1 - x_2)(w) \in M((x_2^{1/T}))[[x_1 - x_2]]$ by Remark 2.6. We can also obtain (2.18) by taking $b = \mathbf{1}$ in (2.15). Taking $l = -1$ in (2.18), we have

$$Y_M^{(s)}(a, \mathbf{1}; i, -1)(w) = \begin{cases} a_i w & \text{if } i \in s/T + \mathbb{Z}, \\ 0 & \text{if } i \notin s/T + \mathbb{Z}. \end{cases} \quad (2.19)$$

By a similar argument, we have $F(\mathbf{1}, a, w|x_1, x_2) = Y_M(a, x_2)w$, $Y_M^{(s)}(\mathbf{1}, a|x_2, x_1 - x_2)(w) = \delta_{s,0} Y_M(a, x_2)w$ and hence

$$Y_M^{(s)}(\mathbf{1}, a; k, l)(w) = \delta_{s,0} \delta_{l,-1} a_k w. \quad (2.20)$$

Lemma 2.9. *Let M be a (V, T) -module. Then, $Y_M(a_{-2}\mathbf{1}, x) = \frac{d}{dx} Y_M(a, x)$.*

Proof. Let $a \in V, w \in M, k \in (1/T)\mathbb{Z}$ and $s \in \mathbb{Z}$ with $0 \leq s \leq T-1$. Taking $j = s/T$ and $l = -2$ in (2.18), we have

$$\begin{aligned} & Y_M^{(s)}(a, \mathbf{1}; \frac{s}{T} + k, -2)w + \frac{s}{T} Y_M^{(s)}(a, \mathbf{1}; \frac{s}{T} + k - 1, -1)w \\ &= \begin{cases} -ka_{s/T+k-1}w & \text{if } k \in \mathbb{Z}, \\ 0 & \text{if } k \notin \mathbb{Z}. \end{cases} \end{aligned} \quad (2.21)$$

Let $r \in \mathbb{Z}$ with $0 \leq r \leq T-1$ and $n \in r/T + \mathbb{Z}$. By (2.19) and (2.21), we have

$$\begin{aligned} (a_{-2}\mathbf{1})_n w &= \sum_{s=0}^{T-1} Y_M^{(s)}(a, \mathbf{1}; n, -2)(w) \\ &= \sum_{s=0}^{T-1} Y_M^{(s)}(a, \mathbf{1}; \frac{s}{T} + (-\frac{s}{T} + n), -2)(w) \\ &= \sum_{s=0}^{T-1} \frac{-s}{T} Y_M^{(s)}(a, \mathbf{1}; \frac{s}{T} + (-\frac{s}{T} + n - 1), -1)(w) - (-\frac{r}{T} + n)a_{n-1}w \\ &= \frac{-r}{T}a_{n-1}w - (-\frac{r}{T} + n)a_{n-1}w \\ &= -na_{n-1}w. \end{aligned}$$

□

3 Subspaces of $\mathbb{C}[z, z^{-1}]$

Throughout this section we fix a non-positive integer Δ . This is the lowest weight of a graded vertex algebra $V = \oplus_{i=\Delta}^{\infty} V_i$ which will be discussed in Section 4. In this section we introduce a subspace $O_{n,m}^{T,1}(\alpha, \beta; z)$ of $\mathbb{C}[z, z^{-1}]$ (see (3.10) below) for $n, m \in (1/T)\mathbb{N}$ and $\alpha, \beta \in \mathbb{Z}$ and we study its properties. The subspace $O_{n,m}^{T,1}(\alpha, \beta; z)$ will be used to define the subspace $O_{n,m}^{T,1}(V)$ of V in Section 4.

For $N, q \in \mathbb{Z}$ and $Q \in \mathbb{Q}$, $O(N, Q, q; z)$ denotes the subspace of $\mathbb{C}[z, z^{-1}]$ spanned by

$$\begin{aligned} & \text{Res}_x \left((1+x)^Q x^{q+j} \sum_{i \in \mathbb{Z}_{\leq N}} z^i x^{-i-1} \right) \\ &= \sum_{i=0}^{N-q-j} \binom{Q}{i} z^{i+q+j}, \quad j = 0, -1, \dots \end{aligned} \quad (3.1)$$

and $z^i, i \in \mathbb{Z}_{\geq N+1}$. If $N \leq q$, then $O(N, Q, q; z) = \mathbb{C}[z, z^{-1}]$. A similar computation as in the proof of Lemma 2.8 shows the following lemma (or see [15, Proof of Lemma 2]).

Lemma 3.1. *Fix $N, q \in \mathbb{Z}$, $Q \in \mathbb{Q}$ and $i \in \mathbb{Z}_{\leq N}$. Then*

$$z^i \equiv \sum_{k=1}^{N-q} \sum_{j=1}^k \binom{-Q}{-i+q+j} \binom{Q}{k-j} z^{q+k} \pmod{O(N, Q, q; z)}. \quad (3.2)$$

The proof of the following lemma is similar to that of [15, Lemma 3].

Lemma 3.2. *Let $N \in \mathbb{Z}$, $q_0, \dots, q_{T-1} \in \mathbb{Z}$ and $Q_0, \dots, Q_{T-1} \in \mathbb{Q}$ such that $Q_i \not\equiv Q_j \pmod{\mathbb{Z}}$ for all $i \neq j$. The diagonal map $\mathbb{C}[z, z^{-1}] \ni f \mapsto (f, \dots, f) \in \mathbb{C}[z, z^{-1}]^{\oplus T}$ induces an isomorphism*

$$\mathbb{C}[z, z^{-1}] / \bigcap_{s=0}^{T-1} O(N, Q_s, q_s; z) \rightarrow \bigoplus_{s=0}^{T-1} \mathbb{C}[z, z^{-1}] / O(N, Q_s, q_s; z)$$

as vector spaces.

Proof. It is sufficient to show that the induced map is surjective. Note that $\mathbb{C}[z, z^{-1}]_{\geq N+1}$ is a subspace of $O(N, Q_s, q_s; z)$ for each s . Fix an integer q such that $q \leq \min\{q_0, \dots, q_{T-1}\}$. We may assume $q \leq N$ from the comment right after (3.1). Since $O(N, Q_s, q; z)$ is a subspace of $O(N, Q_s, q_s; z)$ for each $s = 0, \dots, T-1$, it is sufficient to show that the diagonal map

$$\begin{aligned} & \mathbb{C}[z, z^{-1}]_{N+1-T(N-q), N} \ni f \\ & \mapsto (f + O(N, Q_s, q; z))_{s=0}^{T-1} \in \bigoplus_{s=0}^{T-1} \mathbb{C}[z, z^{-1}] / O(N, Q_s, q; z) \end{aligned}$$

is surjective. For a Laurent polynomial $\Lambda(z) = \sum_{i=N+1-T(N-q)}^N \lambda_i z^i \in \mathbb{C}[z, z^{-1}]_{N+1-T(N-q), N}$, it follows by (3.2) that

$$\Lambda(z) \equiv \sum_{i=N+1-T(N-q)}^N \lambda_i \sum_{k=1}^{N-q} \sum_{j=1}^k \binom{-Q_s}{-i+q+j} \binom{Q_s}{k-j} z^{q+k} \pmod{O(N, Q_s, q; z)} \quad (3.3)$$

for $s = 0, \dots, T-1$. We denote $\sum_{j=1}^k \binom{-Q_s}{-i+q+j} \binom{Q_s}{k-j}$ by $\alpha_{-i+q}^{s,k}$ for $0 \leq s \leq T-1, 1 \leq k \leq N-q$ and $i \in \mathbb{Z}$. Define $T \times T(N-q) \times (N-q)$ -matrices $\Gamma_s, s = 0, \dots, T-1$ by

$$\Gamma_s = \begin{pmatrix} \alpha_{(T-1)(N-q)-1}^{s,1} & \alpha_{(T-1)(N-q)-1}^{s,2} & \cdots & \alpha_{(T-1)(N-q)-1}^{s,N-q} \\ \alpha_{(T-1)(N-q)-2}^{s,1} & \alpha_{(T-1)(N-q)-2}^{s,2} & \cdots & \alpha_{(T-1)(N-q)-2}^{s,N-q} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{-N+q}^{s,1} & \alpha_{-N+q}^{s,2} & \cdots & \alpha_{-N+q}^{s,N-q} \end{pmatrix}.$$

Since

$$(\lambda_{N+1-T(N-q)}, \lambda_{N+2-T(N-q)}, \dots, \lambda_N) \Gamma_s \begin{pmatrix} z^{q+1} \\ \vdots \\ z^N \end{pmatrix}$$

is equal to the right-hand side of (3.3) for $s = 0, \dots, T-1$, it is sufficient to show that the square matrix

$$\Gamma = (\Gamma_0 \ \Gamma_1 \ \cdots \ \Gamma_{T-1}) \quad (3.4)$$

of order $T(N-q)$ is non-singular. It is proved in Subsection 6.1 that Γ is non-singular. $\square$

For $N \in \mathbb{Z}$ and $\gamma \in \mathbb{Q}$, define a linear automorphism $\varphi_{N,\gamma}$ of $\mathbb{C}[z, z^{-1}]$ by

$$\begin{aligned} & \varphi_{N,\gamma}(z^i) \\ &= \begin{cases} (-1)^{i+1} \operatorname{Res}_x \left((1+x)^{\gamma-i} x^i \sum_{j \in \mathbb{Z}_{\leq N}} z^j x^{-j-1} \right) & \text{for } i \leq N, \\ z^i & \text{for } i \geq N+1. \end{cases} \end{aligned} \quad (3.5)$$

Lemma 3.3.

$$\begin{aligned} & \varphi_{N,\gamma}(\operatorname{Res}_x \left((1+x)^k x^i \sum_{j \in \mathbb{Z}_{\leq N}} z^j x^{-j-1} \right)) \\ &= (-1)^{i+1} \operatorname{Res}_x \left((1+x)^{\gamma-k-i} x^i \sum_{j \in \mathbb{Z}_{\leq N}} z^j x^{-j-1} \right) \end{aligned} \quad (3.6)$$

for $k \in \mathbb{Q}$ and $i \in \mathbb{Z}_{\leq N}$. In particular, $\varphi_{N,\gamma}^2 = \operatorname{id}_{\mathbb{C}[z, z^{-1}]}$ and

$$\varphi_{N,\gamma}(O(N, Q, q; z)) = O(N, \gamma - Q - q, q; z)$$

for $Q \in \mathbb{Q}$ and $q \in \mathbb{Z}$.

Proof. We simply write $\varphi = \varphi_{N,\gamma}$. Let $i \in \mathbb{Z}_{\leq N}$. Since

$$\varphi(z^i) = (-1)^{i+1} \sum_{j=0}^{N-i} \binom{\gamma-i}{j} z^{i+j},$$

we have

$$\begin{aligned} \varphi(\text{Res}_x(1+x)^k x^i \sum_{j \in \mathbb{Z}_{\leq N}} z^j x^{-j-1}) &= \sum_{j=0}^{N-i} \binom{k}{j} \varphi(x^{i+j}) \\ &= \sum_{j=0}^{N-i} \binom{k}{j} (-1)^{i+j+1} \sum_{m=0}^{N-i-j} \binom{\gamma-i-j}{m} z^{i+j+m} \\ &= \sum_{j=0}^{N-i} \binom{k}{j} (-1)^{i+j+1+m} \sum_{m=0}^{N-i-j} \binom{-\gamma+i+j+m-1}{m} z^{i+j+m} \\ &= \sum_{l=0}^{N-i} (-1)^{i+1+l} z^{i+l} \sum_{\substack{0 \leq j, m \leq N-i \\ j+m=l}} \binom{k}{j} \binom{-\gamma+i+l-1}{m} \\ &= \sum_{l=0}^{N-i} (-1)^{i+1+l} z^{i+l} \binom{k-\gamma+i+l-1}{l} \\ &= (-1)^{i+1} \sum_{l=0}^{N-i} \binom{-k+\gamma-i}{l} z^{i+l} \\ &= (-1)^{i+1} \text{Res}_x((1+x)^{\gamma-k-i} x^i \sum_{j \in \mathbb{Z}_{\leq N}} z^j x^{-j-1}). \end{aligned}$$

By this, $\varphi^2(z^j) = z^j$ for $j \in \mathbb{Z}$. Since

$$\begin{aligned} \varphi(\text{Res}_x((1+x)^Q x^{q+d} \sum_{j \in \mathbb{Z}_{\leq N}} z^j x^{-j-1})) \\ &= (-1)^{q+d+1} \text{Res}_x((1+x)^{\gamma-Q-q-d} x^{q+d} \sum_{j \in \mathbb{Z}_{\leq N}} z^j x^{-j-1}) \\ &= (-1)^{q+d+1} \sum_{m=0}^{-d} \binom{-d}{m} \text{Res}_x((1+x)^{\gamma-Q-q} x^{q+d+m} \sum_{j \in \mathbb{Z}_{\leq N}} z^j x^{-j-1}) \end{aligned}$$

for $d \in \mathbb{Z}_{\leq 0}$, we have $\varphi_{N,\gamma}(O(N, Q, q; z)) = O(N, \gamma - Q - q, q; z)$. $\square$

Throughout the rest of this section, $m = l_1 + i_1/T, p = l_2 + i_2/T, n = l_3 + i_3/T \in (1/T)\mathbb{N}$ with $l_1, l_2, l_3 \in \mathbb{N}$ and $0 \leq i_1, i_2, i_3 \leq$

$T - 1$. We always denote m, p, n as above until further notice. For $i, j \in (1/T)\mathbb{Z}$, $r(i, j)$ denotes the integer uniquely determined by the conditions

$$0 \leq r(i, j) \leq T - 1 \text{ and } i - j \equiv \frac{r(i, j)}{T} \pmod{\mathbb{Z}}. \quad (3.7)$$

For $0 \leq s \leq T - 1$, $s^\vee$ denotes the integer uniquely determined by the conditions

$$0 \leq s^\vee \leq T - 1 \text{ and } i_1 - i_3 \equiv s + s^\vee \pmod{T}. \quad (3.8)$$

For $s = 0, \dots, T - 1$ and $\alpha, \beta \in \mathbb{Z}$, define

$$\begin{aligned} O_{n,m}^{(T;s),1}(\alpha, \beta; z) &= O(\alpha + \beta - 1 - \Delta, \alpha - 1 + l_1 + \delta(s \leq i_1) + \frac{s}{T}, \\ &\quad - l_1 - l_3 - \delta(s \leq i_1) - \delta(T \leq s + i_3) - 1; z) \end{aligned} \quad (3.9)$$

and

$$O_{n,m}^{T,1}(\alpha, \beta; z) = \bigcap_{s=0}^{T-1} O_{n,m}^{(T;s),1}(\alpha, \beta; z), \quad (3.10)$$

where Δ is the fixed non-positive integer as stated at the beginning of this section and $\delta(i \leq j)$ is defined in (2.1).

For $\alpha, \beta, j \in \mathbb{Z}$ and $s = 0, \dots, T - 1$, define a Laurent polynomial $\Psi_{n,m}^{(T;s)}(\alpha, \beta, j; z) \in \mathbb{C}[z, z^{-1}]$ by

$$\begin{aligned} &\Psi_{n,m}^{(T;s)}(\alpha, \beta, j; z) \\ &= \text{Res}_x \left((1+x)^{\alpha-1+l_1+\delta(s \leq i_1)+s/T} x^{-l_1-l_3-\delta(s \leq i_1)-\delta(T \leq s+i_3)-1+j} \right. \\ &\quad \times \sum_{\substack{i \in \mathbb{Z} \\ i \leq \alpha+\beta-1-\Delta}} z^i x^{-i-1} \Big) \\ &= \sum_{i=0}^{\alpha+\beta-\Delta+l_1+l_3+\delta(s \leq i_1)+\delta(T \leq s+i_3)-j} \binom{\alpha-1+l_1+\delta(s \leq i_1)+\frac{s}{T}}{i} \\ &\quad \times z^{i-l_1-l_3-\delta(s \leq i_1)-\delta(T \leq s+i_3)-1+j}. \end{aligned} \quad (3.11)$$

The disjoint union $\{\Psi_{n,m}^{(T;s)}(\alpha, \beta, j; z) \mid j = 0, -1, \dots\} \cup \{z^i \mid i \geq \alpha + \beta - \Delta\}$ spans $O_{n,m}^{(T;s),1}(\alpha, \beta; z)$ by (3.1).

Lemma 3.4. *Let $m' = l'_1 + i'_1/T, n' = l'_3 + i'_3/T \in (1/T)\mathbb{N}$ with $l'_1, l'_3 \in \mathbb{N}$ and $0 \leq i'_1, i'_3 \leq T-1$. If $m' \leq m$ and $n' \leq n$, then $O_{n,m}^{(T;s),1}(\alpha, \beta; z) \subset O_{n',m'}^{(T;s),1}(\alpha, \beta; z)$ for $\alpha, \beta \in \mathbb{Z}$ and $s = 0, \dots, T-1$. In particular, $O_{n,m}^{T,1}(\alpha, \beta; z) \subset O_{n',m'}^{T,1}(\alpha, \beta; z)$.*

Proof. Let $\rho_1 = l_1 + \delta(s \leq i_1) - (l'_1 + \delta(s \leq i'_1))$ and $\rho_3 = l_3 + \delta(T \leq s + i_3) - (l'_3 + \delta(T \leq s + i'_3))$ for $s = 0, \dots, T-1$. It follows by $m' \leq m$ and $n' \leq n$ that ρ_1 and ρ_3 are non-negative integers. Since

$$\Psi_{n,m}^{(T;s)}(\alpha, \beta, j; z) = \sum_{i=0}^{\rho_1} \binom{\rho_1}{i} \Psi_{n',m'}^{(T;s)}(\alpha, \beta, j - \rho_1 - \rho_3 + i; z),$$

the proof is complete. $\square$

A direct computation shows

$$\frac{-s - s^\vee + i_1 - i_3}{T} + \delta(T \leq s^\vee + i_3) = \delta(s \leq i_1) - 1 \quad (3.12)$$

for $s = 0, \dots, T-1$ and hence

$$\delta(s^\vee \leq i_1) + \delta(T \leq s^\vee + i_3) = \delta(s \leq i_1) + \delta(T \leq s + i_3). \quad (3.13)$$

For a non-positive integer j , it follows by (3.6), (3.12) and (3.13) that

$$\begin{aligned} & \varphi_{\alpha+\beta-1-\Delta, \alpha+\beta+m-n-2}(\Psi_{n,m}^{(T;s^\vee)}(\beta, \alpha, j; z)) \\ &= \text{Res}_x \left((1+x)^{\alpha-1+l_1+\delta(s \leq i_1)+s/T-j} x^{-l_1-l_3-\delta(s \leq i_1)-\delta(T \leq s+i_3)-1+j} \right. \\ & \quad \times \sum_{\substack{i \in \mathbb{Z} \\ i \leq \alpha+\beta-1-\Delta}} z^i x^{-i-1} \Big) \\ &= \sum_{k=0}^{-j} \binom{-j}{k} \Psi_{n,m}^{(T;s)}(\alpha, \beta, j+k; z) \end{aligned}$$

and hence

$$\varphi_{\alpha+\beta-1-\Delta, \alpha+\beta+m-n-2}(O_{n,m}^{(T;s^\vee),1}(\beta, \alpha; z)) = O_{n,m}^{(T;s),1}(\alpha, \beta; z). \quad (3.14)$$

Thus, $\varphi_{\alpha+\beta-1-\Delta, \alpha+\beta+m-n-2}$ induces an isomorphism

$$\mathbb{C}[z, z^{-1}]/O_{n,m}^{(T;s^\vee),1}(\beta, \alpha; z) \rightarrow \mathbb{C}[z, z^{-1}]/O_{n,m}^{(T;s),1}(\alpha, \beta; z) \quad (3.15)$$

and hence

$$\mathbb{C}[z, z^{-1}] / \bigcap_{s=0}^{T-1} O_{n,m}^{(T;s),1}(\beta, \alpha; z) \cong \mathbb{C}[z, z^{-1}] / \bigcap_{s=0}^{T-1} O_{n,m}^{(T;s),1}(\alpha, \beta; z) \quad (3.16)$$

by Lemma 3.1.

Let $\alpha, \beta \in \mathbb{Z}$. Let $q = -l_1 - l_3 - 3$, $Q_{\alpha,s} = \alpha - 1 + l_1 + \delta(s \leq i_1) + s/T$, $s = 0, \dots, T-1$ and

$$R_{\alpha,\beta,\Delta} = \mathbb{C}[z, z^{-1}]_{\alpha+\beta-\Delta-T(\alpha+\beta-1-\Delta-q), \alpha+\beta-1-\Delta}. \quad (3.17)$$

For $r = 0, \dots, T-1$ and $i \in \mathbb{Z}_{\leq \alpha+\beta-1-\Delta}$, it follows by the argument in the proof of Lemma 3.2 that there exists a unique Laurent polynomial in $R_{\alpha,\beta,\Delta}$, which we denote by $E_{n,m}^{(T;r)}(\alpha, \beta, i; z)$, such that

$$\begin{aligned} E_{n,m}^{(T;r)}(\alpha, \beta, i; z) &\in \bigcap_{\substack{0 \leq s \leq T-1 \\ s \neq r}} O(\alpha + \beta - 1 - \Delta, Q_{\alpha,s}, q; z) \quad \text{and} \\ E_{n,m}^{(T;r)}(\alpha, \beta, i; z) - z^i &\in O(\alpha + \beta - 1 - \Delta, Q_{\alpha,r}, q; z). \end{aligned} \quad (3.18)$$

We also define

$$E_{n,m}^{(T;r)}(\alpha, \beta, i; z) = 0 \text{ for } i \in \mathbb{Z}_{\geq \alpha+\beta-\Delta} \quad (3.19)$$

for convenience. Since $O(\alpha + \beta - 1 - \Delta, Q_{\alpha,s}, q; z) \subset O_{n,m}^{(T;s),1}(\alpha, \beta; z)$, we have

$$E_{n,m}^{(T;r)}(\alpha, \beta, i; z) \equiv \delta_{r,s} z^i \pmod{O_{n,m}^{(T;s),1}(\alpha, \beta; z)} \quad (3.20)$$

for $s = 0, \dots, T-1$. Define

$$\begin{aligned} X_{n,m}^{(T;s)}(\alpha, \beta; z, x) &= \sum_{i \in \mathbb{Z}} E_{n,m}^{(T;s)}(\alpha, \beta, i; z) x^{-i-1} \\ &= \sum_{\substack{i \in \mathbb{Z} \\ i \leq \alpha+\beta-1-\Delta}} E_{n,m}^{(T;s)}(\alpha, \beta, i; z) x^{-i-1} \in \mathbb{C}[z, z^{-1}]((x)) \end{aligned} \quad (3.21)$$

for $s = 0, \dots, T-1$. It follows from Lemma 3.2 that

$$\sum_{s=0}^{T-1} E_{n,m}^{(T;s)}(\alpha, \beta, i; z) \equiv z^i \pmod{O_{n,m}^{T,1}(\alpha, \beta; z)} \quad (3.22)$$

for $i \in \mathbb{Z}$ and

$$\sum_{s=0}^{T-1} X_{n,m}^{(T;s)}(\alpha, \beta; z, x) \equiv \sum_{\substack{i \in \mathbb{Z} \\ i \leq \alpha + \beta - 1 - \Delta}} z^i x^{-i-1} \pmod{O_{n,m}^{T,1}(\alpha, \beta; z)((x))}. \quad (3.23)$$

Define a Laurent polynomial $\Phi_{n,p,m}^T(\alpha, \beta; z) \in R_{\alpha,\beta,\Delta}$ by

$$\begin{aligned} & \Phi_{n,p,m}^T(\alpha, \beta; z) \\ &= \sum_{i=0}^{l_2} \binom{-l_1 - l_3 + l_2 - \delta(r(p,n) \leq i_1) - \delta(T \leq r(p,n) + i_3)}{i} \\ & \quad \times \text{Res}_x \left((1+x)^{\alpha-1+l_1+\delta(r(p,n) \leq i_1)+r(p,n)/T} \right. \\ & \quad \times x^{-l_1-l_3+l_2-\delta(r(p,n) \leq i_1)-\delta(T \leq r(p,n)+i_3)-i} X_{n,m}^{(T;r(p,n))}(\alpha, \beta; z, x) \Big), \end{aligned} \quad (3.24)$$

where $r(p, n)$ is defined in (3.7). This is used to define the product $*_{n,p,m}^T$ on a vertex algebra in Section 4.

We denote $\text{Span}_{\mathbb{C}}\{z^i \in \mathbb{C}[z, z^{-1}] \mid i \neq -1\}$ by $\mathbb{C}[z, z^{-1}]_{\neq -1}$. The following two results will be used to compute $\mathbf{1} *_{n,p,m}^T a$ for $a \in V$ in Section 4.

Lemma 3.5. *Let $\alpha, r \in \mathbb{Z}$ with $0 \leq r \leq T-1$. Then*

$$X_{n,m}^{(T;r)}(0, \alpha; z, x) \equiv \delta_{r,0} z^{-1} \pmod{(\mathbb{C}[z, z^{-1}]_{\neq -1}((x)))}.$$

Proof. Since

$$\Psi_{n,m}^{(T;0)}(0, \alpha, j; z) = \sum_{i=0}^{\alpha+\Delta+l_1+l_3+1-j} \binom{l_1}{i} z^{i-l_1-l_3-2+j} \in \mathbb{C}[z, z^{-1}]_{\leq -2}$$

for all $j \in \mathbb{Z}_{\leq 0}$, $O_{n,m}^{(T;0),1}(0, \alpha; z)$ is a subspace of $\mathbb{C}[z, z^{-1}]_{\neq -1}$. Since $E_{n,m}^{(T;r)}(\alpha, \beta, i; z) - \delta_{r,0} z^i \in O_{n,m}^{(T;0),1}(0, \alpha; z)$ for $i \in \mathbb{Z}$, the proof is complete. $\square$

Lemma 3.6. *Let $\alpha \in \mathbb{Z}$. Then*

$$\Phi_{n,p,m}^T(0, \alpha; z) \equiv \delta_{n,p} z^{-1} \pmod{\mathbb{C}[z, z^{-1}]_{\neq -1}}.$$

Proof. If $n \not\equiv p \pmod{\mathbb{Z}}$, then it follows by Lemma 3.5 that

$$\Phi_{n,p,m}^T(0, \alpha; z) \equiv 0 \pmod{\mathbb{C}[z, z^{-1}]_{\neq -1}}.$$

Suppose $n \equiv p \pmod{\mathbb{Z}}$. By Lemma 3.5 again, the same computation as in the proof of [4, Lemma 4.7] shows

$$\Phi_{n,p,m}^T(0, \alpha; z) \equiv \delta_{n,p} z^{-1} \pmod{\mathbb{C}[z, z^{-1}]_{\neq -1}}.$$

□

The following result will be used in order to obtain Lemma 4.3, which induces the commutator formula in Lemma 5.9.

Lemma 3.7. *Let $\alpha, \beta \in \mathbb{Z}$.*

$$\begin{aligned} & \Phi_{n,p,m}^T(\alpha, \beta; z) - \varphi_{\alpha+\beta-1-\Delta, \alpha+\beta+m-n-2}(\Phi_{n,m+n-p,m}^T(\beta, \alpha; z)) \\ & - \text{Res}_x(1+x)^{\alpha-1+p-n} X_{n,m}^{(T;r(p,n))}(\alpha, \beta; z, x) \in O_{n,m}^{T,1}(\alpha, \beta; z). \end{aligned} \quad (3.25)$$

Proof. The proof is similar to that of [5, Lemma 3.4]. We simply write $r = r(p, n)$ and $\varphi = \varphi_{\alpha+\beta-1-\Delta, \alpha+\beta+m-n-2}$. It follows by

$$(m+n-p) - n \equiv \frac{i_1 - i_2}{T} \equiv \frac{r^\vee}{T} \pmod{\mathbb{Z}}$$

that $\Phi_{n,m+n-p,p}^T(\beta, \alpha; z) \in \cap_{s \neq r^\vee} O_{n,m}^{(T;s),1}(\beta, \alpha; z)$, where $r^\vee$ is defined in (3.8). Since $\varphi(\Phi_{n,m+n-p,p}^T(\beta, \alpha; z)) \in \cap_{s \neq r} O_{n,m}^{(T;s),1}(\alpha, \beta; z)$ by (3.14), we have

$$\begin{aligned} & \Phi_{n,p,m}^T(\alpha, \beta; z) - \varphi(\Phi_{n,m+n-p,m}^T(\beta, \alpha; z)) \\ & - \text{Res}_x(1+x)^{\alpha-1+p-n} X_{n,m}^{(T;r)}(\alpha, \beta; z, x) \in \bigcap_{s \neq r} O_{n,m}^{(T;s),1}(\alpha, \beta; z). \end{aligned}$$

Thus, it is sufficient to show (3.25) modulo $O_{n,m}^{(T;r),1}(\alpha, \beta; z)$ by Lemma 3.2. Define

$$\varepsilon = \begin{cases} 1 & \text{if } T \leq i_1 + i_3 - i_2, \\ 0 & \text{if } 0 \leq i_1 + i_3 - i_2 < T, \\ -1 & \text{if } i_1 + i_3 - i_2 < 0. \end{cases} \quad (3.26)$$

It follows by the formula of ε in the proof of [5, Lemma 3.4] and (3.13) that

$$\begin{aligned}
& \Phi_{n,m+n-p,m}^T(\beta, \alpha; z) \\
&= \sum_{i=0}^{l_1+l_3-l_2+\varepsilon} \binom{-l_1-l_3+(l_1+l_3-l_2+\varepsilon)-\delta(r^\vee \leq i_1)-\delta(T \leq r^\vee+i_3)}{i} \\
&\quad \times \text{Res}_x(1+x)^{\beta-1+l_1+\delta(r^\vee \leq i_1)+r^\vee/T} \\
&\quad \times x^{-l_1-l_3+(l_1+l_3-l_2+\varepsilon)-\delta(r^\vee \leq i_1)-\delta(T \leq r^\vee+i_3)-i} X_{n,m}^{(T;r^\vee)}(\beta, \alpha; z, x) \\
&= \sum_{i=0}^{l_1+l_3-l_2+\varepsilon} \binom{-l_2-1}{i} \text{Res}_x(1+x)^{\beta-1+l_1+\delta(r^\vee \leq i_1)+r^\vee/T} x^{-l_2-1-i} X_{n,m}^{(T;r^\vee)}(\beta, \alpha; z, x).
\end{aligned}$$

Thus, it follows by (3.6) that

$$\begin{aligned}
& \varphi(\Phi_{n,m+n-p,m}^T(\beta, \alpha; z)) \\
&= \sum_{i=0}^{l_1+l_3-l_2+\varepsilon} \binom{-l_2-1}{i} \text{Res}_x(1+x)^{\alpha-1+p-n+i} x^{-l_2-1-i} X_{n,m}^{(T;r)}(\alpha, \beta; z, x) \\
&\hspace{15em} (3.27)
\end{aligned}$$

and therefore

$$\begin{aligned}
& \varphi(\Phi_{n,m+n-p,m}^T(\beta, \alpha; z)) \\
&\equiv \sum_{i=0}^{l_1+l_3-l_2+\varepsilon} \binom{-l_2-1}{i} \text{Res}_x(1+x)^{\alpha-1+p-n+i} x^{-l_2-1-i} \sum_{\substack{j \in \mathbb{Z} \\ j \leq \alpha+\beta-1-\Delta}} z^j x^{-j-1} \\
&\quad (\text{mod } O_{n,m}^{(T;r),1}(\alpha, \beta; z)).
\end{aligned}$$

The same argument as in the proof of [5, Lemma 3.4] shows

$$\begin{aligned}
& \Phi_{n,p,m}^T(\alpha, \beta; z) - \varphi(\Phi_{n,m+n-p,m}^T(\beta, \alpha; z)) \\
& \equiv \sum_{i=0}^{l_2} \binom{-l_1 - l_3 + l_2 - \varepsilon - 1}{i} \\
& \quad \times \operatorname{Res}_x \left((1+x)^{\alpha-1+l_1+\delta(r \leq i_1)+r/T} x^{-l_1-l_3+l_2-\varepsilon-1-i} \sum_{\substack{j \in \mathbb{Z} \\ j \leq \alpha+\beta-1-\Delta}} z^j x^{-j-1} \right) \\
& \quad - \sum_{i=0}^{l_1+l_3-l_2+\varepsilon} \binom{-l_2-1}{i} \operatorname{Res}_x (1+x)^{\alpha-1+p-n+i} x^{-l_2-1-i} \sum_{\substack{j \in \mathbb{Z} \\ j \leq \alpha+\beta-1-\Delta}} z^j x^{-j-1} \\
& \quad (\text{mod } O_{n,m}^{(T;r),1}(\alpha, \beta; z)) \\
& = \operatorname{Res}_x (1+x)^{\alpha-1+p-n} \sum_{\substack{j \in \mathbb{Z} \\ j \leq \alpha+\beta-1-\Delta}} z^j x^{-j-1}.
\end{aligned}$$

The proof is complete. $\square$

Let $l \in (1/T)\mathbb{N}$ with $l \leq n, m$. Then, it follows by Lemma 3.4 that $X_{n,m}^{(T;r)}(\alpha, \beta; z, x) \equiv X_{n-l,m-l}^{(T;r)}(\alpha, \beta; z, x) \pmod{O_{n-l,m-l}^{T,1}(\alpha, \beta; z)((x))}$ for $\alpha, \beta \in \mathbb{Z}$. The same computation as in the proof of [5, Proposition 4.3] shows the following lemma.

Lemma 3.8. *Let $l \in (1/T)\mathbb{N}$ with $l \leq n, m$. Then*

$$\Phi_{n,p,m}^T(\alpha, \beta; z) \equiv \Phi_{n-l,p-l,m-l}^T(\alpha, \beta; z) \pmod{O_{n-l,m-l}^{T,1}(\alpha, \beta; z)}$$

for $\alpha, \beta \in \mathbb{Z}$.

Let T' be a positive multiple of T and $\alpha, \beta \in \mathbb{Z}$. Set $d = T'/T$. We note that $m = l_1 + di_1/T', p = l_2 + di_2/T'$ and $n = l_3 + di_3/T'$. Thus it follows by (3.9) that

$$O_{n,m}^{(T';dr),1}(\alpha, \beta; z) = O_{n,m}^{(T;r),1}(\alpha, \beta; z) \quad (3.28)$$

for $r = 0, \dots, T-1$. By this and (3.20), we have

$$E_{n,m}^{(T';dr)}(\alpha, \beta, i; z) \equiv \delta_{r,s} z^i \pmod{O_{n,m}^{(T;s),1}(\alpha, \beta, i; z)}$$

for $i \in \mathbb{Z}$ and $r, s = 0, \dots, T-1$. Therefore, Lemma 3.2 implies

$$E_{n,m}^{(T';dr)}(\alpha, \beta, i; z) \equiv E_{n,m}^{(T;r)}(\alpha, \beta, i; z) \pmod{O_{n,m}^{T,1}(\alpha, \beta; z)}$$

for $i \in \mathbb{Z}$ and $r = 0, \dots, T-1$. By (3.21) and (3.24), we have the following result.

Lemma 3.9. *Let T' be a positive multiple of T and $\alpha, \beta \in \mathbb{Z}$. Then*

$$\Phi_{n,p,m}^{T'}(\alpha, \beta; z) \equiv \Phi_{n,p,m}^T(\alpha, \beta; z) \pmod{O_{n,m}^{T,1}(\alpha, \beta; z)}.$$

4 Associative algebras $A_m^T(V)$ and bi-modules $A_{n,m}^T(V)$

Throughout the rest of this paper, we always assume the following properties for a vertex algebra V : V has a grading $V = \bigoplus_{i=\Delta}^{\infty} V_i$ such that $\Delta \in \mathbb{Z}_{\leq 0}$, $\mathbf{1} \in V_0$ and for all homogeneous element $a \in V$, $a_i V_j \subset V_{\text{wt } a - 1 - i + j}$, where $V_i = 0$ for $i < \Delta$. Every vertex operator algebra satisfies these properties. Throughout this section, we fix $m = l_1 + i_1/T, p = l_2 + i_2/T, n = l_3 + i_3/T \in (1/T)\mathbb{N}$ with $l_1, l_2, l_3 \in \mathbb{N}$ and $0 \leq i_1, i_2, i_3 \leq T - 1$.

In this section, we first define a product $*_{n,p,m}^T$ on V and a quotient space $A_{n,m}^T(V)$ of V . In the following, we shall use a similar argument as in [4, Section 3]. For $a \in V_i$, we denote i by $\text{wt } a$. Define

$$\hat{E}_{n,m}^{(T;s)}(a, b, i) = E_{n,m}^{(T;s)}(\text{wt } a, \text{wt } b, i; z)|_{z^j = a_j b} \in V \quad (4.1)$$

for homogeneous elements a, b of V and $i \in \mathbb{Z}$ and extend $\hat{E}_{n,m}^{(T;s)}(a, b, i)$ for arbitrary $a, b \in V$ by linearity.

Let $O_{n,m}^{T,0}(V)$ be the subspace of V spanned by

$$\{a_{-2}\mathbf{1} + (\text{wt } a + m - n)a \in V \mid \text{homogeneous } a \in V\} \quad (4.2)$$

and $O_{n,m}^{T,1}(V)$ the subspace of V spanned by

$$\left\{ P(z)|_{z^j = a_j b} \in V \mid \begin{array}{l} \text{homogeneous } a, b \in V \text{ and} \\ P(z) \in O_{n,m}^{T,1}(\text{wt } a, \text{wt } b; z) \end{array} \right\}. \quad (4.3)$$

A similar argument as in the proof of [16, Lemma 2.1.3] shows the following lemma as stated in the proof of [4, Lemm 2.3].

Lemma 4.1.

$$\begin{aligned} & \text{Res}_x(1+x)^i x^j Y(b, x)a \\ & \equiv (-1)^{j+1} \text{Res}_x(1+x)^{\text{wt } a + \text{wt } b + m - n - 2 - i - j} x^j Y(a, x)b \pmod{O_{n,m}^{T,0}(V)}. \end{aligned}$$

for $i \in \mathbb{Q}, j \in \mathbb{Z}$ and homogeneous $a, b \in V$.

Define

$$\begin{aligned}\hat{X}_{n,m}^{(T;s)}(a, b; x) &= X_{n,m}^{(T;s)}(\text{wt } a, \text{wt } b; z, x)|_{z^j=a_j b} \\ &= \sum_{\substack{i \in \mathbb{Z} \\ i \leq \text{wt } a + \text{wt } b - 1 - \Delta}} \hat{E}_{n,m}^{(T;s)}(a, b, i) x^{-i-1}\end{aligned}\quad (4.4)$$

for $a, b \in V$ and $s = 0, \dots, T-1$. By (3.23), we have

$$\sum_{s=0}^{T-1} \hat{E}_{n,m}^{(T;s)}(a, b, i) \equiv a_i b \pmod{O_{n,m}^{T,1}(V)} \quad (4.5)$$

for $i \in \mathbb{Z}$ and

$$\sum_{s=0}^{T-1} \hat{X}_{n,m}^{(T;s)}(a, b; x) \equiv Y(a, x)b \pmod{O_{n,m}^{T,1}(V)((x))}. \quad (4.6)$$

Define

$$a *_{n,p,m}^T b = \Phi_{n,p,m}^T(\text{wt } a, \text{wt } b; z)|_{z^j=a_j b} \in V \quad (4.7)$$

for homogeneous $a, b \in V$ and extend $a *_{n,p,m}^T b$ for arbitrary $a, b \in V$ by linearity. By $Y(\mathbf{1}, x) = \text{id}_V$ and Lemma 3.6, we have

$$\mathbf{1} *_{n,p,m}^T a = \delta_{n,p} a \quad (4.8)$$

for $a \in V$.

Definition 4.2. Let $O_{n,m}^{T,2}(V)$ be the subspace of V spanned by

$$u *_{n,p_3,m}^T ((a *_{p_3,p_2,p_1}^T b) *_{p_3,p_1,m}^T c - a *_{p_3,p_2,m}^T (b *_{p_2,p_1,m}^T c))$$

for all $a, b, c, u \in V$ and all $p_1, p_2, p_3 \in (1/T)\mathbb{N}$. Define

$$\begin{aligned}O_{n,m}^{T,3}(V) &= \sum_{p_1, p_2 \in (1/T)\mathbb{N}} (V *_{n,p_2,p_1}^T (O_{p_2,p_1}^{T,0}(V) + O_{p_2,p_1}^{T,1}(V)) *_{n,p_1,m}^T V\end{aligned}$$

and

$$O_{n,m}^T(V) = O_{n,m}^{T,0}(V) + O_{n,m}^{T,1}(V) + O_{n,m}^{T,2}(V) + O_{n,m}^{T,3}(V).$$

By (4.8), we have

$$(a *_{n,p_2,p_1}^T b) *_{n,p_1,m}^T c - a *_{n,p_2,m}^T (b *_{p_2,p_1,m}^T c) \in O_{n,m}^{T,2}(V)$$

for $a, b, c \in V$ and $p_1, p_2 \in (1/T)\mathbb{N}$.

Lemma 4.3.

$$\begin{aligned} & a *_{n,p,m}^T b - b *_{n,m+n-p,m}^T a - \text{Res}_x(1+x)^{\text{wt } a-1+p-n} \hat{X}_{n,m}^{(T;r(p,n))}(a, b; x) \\ & \in O_{n,m}^{T,0}(V) + O_{n,m}^{T,1}(V), \end{aligned}$$

where $r(p, n)$ is defined in (3.7).

Proof. We simply write $r = r(p, n)$. Let ε be the integer defined in (3.26). By Lemma 4.1 and (3.27), we have

$$\begin{aligned} & b *_{n,m+p-n,m}^T a \\ & \equiv \sum_{i=0}^{l_1+l_3-l_2+\varepsilon} \binom{-l_2-1}{i} \text{Res}_x(1+x)^{\text{wt } a-1+p-n+i} x^{-l_2-1-i} \hat{X}_{n,m}^{(T;r)}(a, b; x) \\ & \quad (\text{mod } O_{n,m}^{T,0}(V) + O_{n,m}^{T,1}(V)) \\ & = \sum_{i=0}^{l_1+l_3-l_2+\varepsilon} \binom{-l_2-1}{i} \text{Res}_x(1+x)^{\text{wt } a-1+p-n+i} x^{-l_2-1-i} \\ & \quad \times X_{n,m}^{(T;r)}(\text{wt } a, \text{wt } b; z, x)|_{z^j=a_j b} \\ & = \varphi_{\text{wt } a+\text{wt } b-1-\Delta, \text{wt } a+\text{wt } b+m-n-2}(\Phi_{n,m+n-p,m}^T(\text{wt } b, \text{wt } a; z))|_{z^j=a_j b}, \end{aligned}$$

where $\varphi_{\text{wt } a+\text{wt } b-1-\Delta, \text{wt } a+\text{wt } b+m-n-2}$ is defined by (3.5). Thus, the assertion follows from Lemma 3.7. $\square$

By (4.8) and Lemmas 3.5 and 4.3, we have

$$a *_{n,m,m}^T \mathbf{1} \equiv a \pmod{O_{n,m}^{T,0}(V) + O_{n,m}^{T,1}(V)} \quad (4.9)$$

for $a \in V$.

The same argument as in the proof of [5, Lemma 3.8] shows the following lemma.

Lemma 4.4. For $m, p, n \in (1/T)\mathbb{Z}$, we have $V *_{n,p,m}^T O_{p,m}^T(V) \subset O_{n,m}^T(V)$ and $O_{n,p}^T(V) *_{n,p,m}^T V \subset O_{n,m}^T(V)$.

We define

$$A_{n,m}^T(V) = V/O_{n,m}^T(V). \quad (4.10)$$

If $m = n$, we simply write $A_m^T(V) = A_{m,m}^T(V)$. By Definition 4.2, (4.8), (4.9) and Lemma 4.4, we have the following result.

Theorem 4.5. *Let $m, n \in (1/T)\mathbb{N}$. Then, $(A_m^T(V), *_{n,m,m}^T)$ is an associative $\mathbb{C}$ -algebra and $A_{n,m}^T(V)$ is an $A_n^T(V)$ - $A_m^T(V)$ -bimodule, where the left action of $A_n^T(V)$ is given by $*_{n,n,m}^T$ and the right action of $A_m^T(V)$ is given by $*_{n,m,m}^T$.*

Lemmas 3.4 and 3.8 imply the following result.

Proposition 4.6. *Let $l, m, n \in (1/T)\mathbb{N}$ with $l \leq n, m$. Then $O_{n,m}^{T,1}(V)$ is a subspace of $O_{n-l,m-l}^{T,1}(V)$. Moreover, the identity map on V induces a surjective algebra homomorphism $A_m^T(V) \rightarrow A_{m-l}^T(V)$ and a surjective $A_n^T(V)$ - $A_m^T(V)$ -bimodule homomorphism $A_{n,m}^T(V) \rightarrow A_{n-l,m-l}^T(V)$.*

Lemma 3.9 and (3.28) imply the following result.

Proposition 4.7. *Let $m, n \in (1/T)\mathbb{N}$ and T' a positive multiple of T . Then $O_{n,m}^{T,1}(V)$ is a subspace of $O_{n,m}^{T',1}(V)$. Moreover, the identity map on V induces a surjective algebra homomorphism $A_m^{T'}(V) \rightarrow A_m^T(V)$ and a surjective $A_n^{T'}(V)$ - $A_m^{T'}(V)$ -bimodule homomorphism $A_{n,m}^{T'}(V) \rightarrow A_{n,m}^T(V)$.*

Remark 4.8. Suppose V is a vertex operator algebra. Let g be an automorphism of V of finite order t . In [5], a product $*_{g,m,p}^n$ on V and a quotient space $A_{g,n,m}(V) = V/O_{g,n,m}(V)$ of V are constructed for each $n, p, m \in (1/t)\mathbb{N}$. If $g = \text{id}_V$, then $*_{g,m,p}^n = *_{m,p}^n$ and $A_{g,n,m}(V) = A_{n,m}(V)$, where $*_{m,p}^n$ is a product on V and $A_{n,m}(V)$ is a quotient space of V constructed in [4].

We shall discuss a relation between $A_{g,n,m}(V)$ and $A_{n,m}^T(V)$. Suppose $T = 1$. Then $*_{n,p,m}^1 = *_{m,p}^n$ by the definition. Moreover, $O_{n,m}^{1,0}(V) + O_{n,m}^{1,1}(V) = O'_{n,m}(V)$ by (3.9) and (3.10), where $O'_{n,m}(V)$ is the subspace of V defined on p. 801 in [4]. Thus, $O_{n,m}^1(V) = O_{n,m}(V)$ and $A_{n,m}^1(V) = A_{n,m}(V)$.

We shall use the notation in Remark 2.7 and [5]. For homogeneous $a, b \in V$ and $P(z) \in O_{n,m}^{t,1}(\text{wt } a, \text{wt } b; z)$, the definition of $O_{n,m}^{t,1}(\text{wt } a, \text{wt } b; z)$ implies

$$P(z)|_{z^j=a_j b} = \sum_{r=0}^{t-1} P(z)|_{z^j=a_j^{(g,r)} b} \in O'_{g,n,m}(V),$$

where $O'_{g,n,m}(V)$ is the subspace of V defined on p. 4240 in [5]. Thus, $O_{n,m}^{t,1}(V)$ is a subspace of $O'_{g,n,m}(V)$. We simply write $r = r(p, n)$, which is defined in (3.7). For $s = 0, \dots, t-1$, we have

$$\begin{aligned} & \hat{E}_{n,m}^{(t;r)}(a^{(g,s)}, b, i) - \delta_{r,s} a_i^{(g,s)} b \\ &= (E_{n,m}^{(t;r)}(\text{wt } a^{(g,s)}, \text{wt } b, i; z) - \delta_{r,s} z^i)|_{z^j = a_j^{(g,s)} b} \in O'_{g,n,m}(V) \end{aligned}$$

since $E_{n,m}^{(t;r)}(\text{wt } a^{(g,s)}, \text{wt } b, i; z) - \delta_{r,s} z^i \in O_{n,m}^{(t;s),1}(\text{wt } a^{(g,s)}, \text{wt } b; z)$ by (3.20). Therefore, by (3.24) and (4.7) we have

$$\begin{aligned} a *_{n,p,m}^t b &= \sum_{s \neq r} a^{(g,s)} *_{n,p,m}^t b + a^{(g,r)} *_{n,p,m}^t b \\ &\equiv a^{(g,r)} *_{g,m,p}^n b \pmod{O'_{g,n,m}(V)}. \end{aligned}$$

We conclude that $O_{n,m}^{t,1}(V) \subset O_{g,n,m}(V)$ and $A_{g,n,m}(V)$ is a quotient space of $A_{n,m}^t(V)$.

For an automorphism group G of V of finite order, the same argument as above shows $A_{G,n}(V)$ in [15] is a quotient space of $A_n^{|G|}(V)$.

5 $(1/T)\mathbb{N}$ -graded (V, T) -modules and $A_{n,m}^T(V)$

Throughout this section, we always assume the properties mentioned at the beginning of Section 4 for a vertex algebra V as stated there. In this section, for $m \in (1/T)\mathbb{N}$ we describe a relation between the $A_m^T(V)$ -modules and the $(1/T)\mathbb{N}$ -graded (V, T) -modules defined below.

Definition 5.1. A $(1/T)\mathbb{N}$ -graded (V, T) -module M is a (V, T) -module with a $(1/T)\mathbb{N}$ -grading $M = \oplus_{n \in (1/T)\mathbb{N}} M(n)$ such that

$$a_i M(n) \subset M(n + \text{wt } a - i - 1)$$

for homogeneous $a \in V$ and $i, n \in (1/T)\mathbb{N}$.

For a $(1/T)\mathbb{N}$ -graded (V, T) -module M , a (V, T) -submodule N of M is called $(1/T)\mathbb{N}$ -graded (V, T) -submodule of M if N is a $(1/T)\mathbb{N}$ -graded (V, T) -module such that every homogeneous subspace of N is contained in some homogeneous subspace of M . A non-zero $(1/T)\mathbb{N}$ -graded (V, T) -module M is called *simple* if there is no $(1/T)\mathbb{N}$ -graded submodules of M except 0 and M itself.

In the following, we shall use a similar argument as in [4, Section 4]. Throughout this section, $m = l_1 + i_1/T, n = l_3 + i_3/T \in (1/T)\mathbb{N}$ with $l_1, l_2 \in \mathbb{N}$ and $0 \leq i_1, i_3 \leq T-1$. Until Proposition 5.7, $M = \oplus_{i \in (1/T)\mathbb{N}} M(i)$ is a $(1/T)\mathbb{N}$ -graded (V, T) -module. Without loss of generality, we can shift the grading of a $(1/T)\mathbb{N}$ -graded (V, T) -module M so that $M(0) \neq 0$ if $M \neq 0$.

Define a linear map $o_{n,m} : V \rightarrow \text{Hom}_{\mathbb{C}}(M(m), M(n))$ by

$$o_{n,m}(a) = a_{\text{wt } a + m - n - 1} \quad (5.1)$$

for homogeneous $a \in V$ and extend $o_{n,m}(a)$ for arbitrary $a \in V$ by linearity. If $m = n$, we simply write $o = o_{m,m}$. Define a linear map $Z_{M,n,m}^{(s)}(a, b; -) : \mathbb{C}[z, z^{-1}] \rightarrow \text{Hom}_{\mathbb{C}}(M(m), M)$ by

$$Z_{M,n,m}^{(s)}(a, b; z^i) = Y_M^{(s)}(a, b; \text{wt } a + \text{wt } b + m - n - 2 - i, i) \quad (5.2)$$

for $s = 0, \dots, T-1$ and homogeneous $a, b \in V$ and extend $Z_{M,n,m}^{(s)}(a, b; -)$ for arbitrary elements $a, b \in V$ by linearity. Lemma 2.8 implies that the image of $Z_{M,n,m}^{(s)}(a, b; f(z)) : M(m) \rightarrow M$ is contained in $M(n)$ for $f(z) \in \mathbb{C}[z, z^{-1}]$. That is, $Z_{M,n,m}^{(s)}(a, b; -) : \mathbb{C}[z, z^{-1}] \rightarrow \text{Hom}_{\mathbb{C}}(M(m), M(n))$.

Lemma 5.2. *For $s = 0, \dots, T-1$ and homogeneous $a, b \in V$, $Z_{M,n,m}^{(s)}(a, b; -) = 0$ on $O_{n,m}^{(T;s),1}(\text{wt } a, \text{wt } b; z)$.*

Proof. It is sufficient to show that $Z_{M,n,m}^{(s)}(a, b; \Psi_{n,m}^{(T;s)}(\text{wt } a, \text{wt } b, d; z)) = 0$ for all $d \in \mathbb{Z}_{\leq 0}$. Let $w \in M(m)$. Since $Y_M(Y(a, x_0)b, x_2)w \in M((x_2^{1/T})((x_0))_{\geq -\text{wt } a - \text{wt } b + \Delta})$, it follows by Remark 2.6 that

$$Y_M^{(s)}(a, b|x_2, x_0)(w) \in M((x_2^{1/T})((x_0))_{\geq -\text{wt } a - \text{wt } b + \Delta}). \quad (5.3)$$

Let

$$\begin{aligned} j &= \text{wt } a - 1 + l_1 + \delta(s \leq i_1) + \frac{s}{T}, \\ k &= \text{wt } b - 1 + l_1 + \delta(s^\vee \leq i_1) + \frac{s^\vee}{T} - d \quad \text{and} \\ l &= -l_1 - l_3 - \delta(s \leq i_1) - \delta(T \leq s + i_3) - 1 + d, \end{aligned}$$

where $s^\vee$ is defined in (3.8). Since $a_{j+i} = b_{k+i} = 0$ on $M(m)$ for all $i \in \mathbb{N}$, it follows by (2.15), (3.12) and (5.3) that

$$\begin{aligned}
& Z_{M,n,m}^{(s)}(a, b; \Psi_{n,m}^{(T;s)}(\text{wt } a, \text{wt } b, d; z))(w) \\
&= \sum_{i=0}^{\text{wt } a + \text{wt } b - 1 - \Delta - l} \binom{j}{i} Y_M^{(s)}(a, b; j + k - i, l + i)(w) \\
&= \sum_{i=0}^{\infty} \binom{j}{i} Y_M^{(s)}(a, b; j + k - i, l + i)(w) \\
&= 0.
\end{aligned}$$

□

Lemma 5.3. For $u \in O_{n,m}^{T,0}(V) + O_{n,m}^{T,1}(V)$, $o_{n,m}(u) = 0$ on $M(m)$.

Proof. Let a, b be homogeneous elements of V . It follows by Lemma 2.9 that $o_{n,m}(a_{-2}\mathbf{1} + (\text{wt } a + m - n)a) = 0$ on $M(m)$. Let $P(z) = \sum_{i \in \mathbb{Z}} \lambda_i z^i \in O_{n,m}^{T,1}(\text{wt } a, \text{wt } b; z)$. It follows by Lemma 5.2 that on $M(m)$

$$\begin{aligned}
o_{n,m}\left(\sum_{i \in \mathbb{Z}} \lambda_i a_i b\right) &= \sum_{i \in \mathbb{Z}} \lambda_i o_{n,m}(a_i b) \\
&= \sum_{s=0}^{T-1} \sum_{i \in \mathbb{Z}} \lambda_i Y_M^{(s)}(a, b; \text{wt } a + \text{wt } b + m - n - 2 - i, i) \\
&= \sum_{s=0}^{T-1} \sum_{i \in \mathbb{Z}} \lambda_i Z_{M,n,m}^{(s)}(a, b; z^i) \\
&= \sum_{s=0}^{T-1} Z_{M,n,m}^{(s)}(a, b; P(z)) \\
&= 0.
\end{aligned}$$

□

Lemma 5.4. For $a, b \in V$ and $w \in M(m)$

$$o_{n,m}(a *_{n,p,m}^T b)w = o_{n,p}(a) o_{p,m}(b)w.$$

Proof. We may assume a and b to be homogeneous elements of V . We simply write $r = r(p, n)$, which is defined in (3.7). By (3.20) and

Lemma 5.2, we have

$$\begin{aligned}
& Z_{M,n,m}^{(r)}(a, b; E_{n,m}^{(T;r)}(\text{wt } a, \text{wt } b, j; z))(w) \\
&= Z_{M,n,m}^{(r)}(a, b; z^j)(w) \\
&= Y_M^{(r)}(a, b; \text{wt } a + \text{wt } b + m - n - 2 - j, j) \quad (5.4)
\end{aligned}$$

for $j \in \mathbb{Z}$. We write $\Phi_{n,p,m}^T(\text{wt } a, \text{wt } b; z) = \sum_{i \in \mathbb{Z}} \lambda_i z^i, \lambda_i \in \mathbb{C}$. By (5.4), we have

$$\begin{aligned}
o_{n,m}(a *_{n,p,m}^T b)w &= \sum_{i \in \mathbb{Z}} \lambda_i o_{n,m}(a_i b)w \\
&= \sum_{s=0}^{T-1} \sum_{i \in \mathbb{Z}} \lambda_i Y_M^{(s)}(a, b; \text{wt } a + \text{wt } b + m - n - 2 - i, i)(w) \\
&= \sum_{s=0}^{T-1} Z_{M,n,m}^{(s)}(a, b; \Phi_{n,p,m}^T(\text{wt } a, \text{wt } b; z))(w) \\
&= Z_{M,n,m}^{(r)}(a, b; \Phi_{n,p,m}^T(\text{wt } a, \text{wt } b; z))(w) \\
&= \sum_{i=0}^{l_2} \binom{-l_1 - l_3 + l_2 - \delta(r \leq i_1) - \delta(T \leq r + i_3)}{i} \\
&\quad \times \text{Res}_x \left((1+x)^{\text{wt } a - 1 + l_1 + \delta(r \leq i_1) + r/T} x^{-l_1 - l_3 + l_2 - \delta(r \leq i_1) - \delta(T \leq r + i_3) - i} \right. \\
&\quad \times \sum_{j \in \mathbb{Z}} Z_{M,n,m}^{(r)}(a, b; E_{n,m}^{(T;r)}(\text{wt } a, \text{wt } b, j; z))(w) x^{-j-1} \Big) \\
&= \sum_{i=0}^{l_2} \binom{-l_1 - l_3 + l_2 - \delta(r \leq i_1) - \delta(T \leq r + i_3)}{i} \\
&\quad \times \text{Res}_x \left((1+x)^{\text{wt } a - 1 + l_1 + \delta(r \leq i_1) + r/T} x^{-l_1 - l_3 + l_2 - \delta(r \leq i_1) - \delta(T \leq r + i_3) - i} \right. \\
&\quad \times \sum_{j \in \mathbb{Z}} Y_M^{(r)}(a, b; \text{wt } a + \text{wt } b + m - n - 2 - j, j)(w) x^{-j-1} \Big). \quad (5.5)
\end{aligned}$$

Let $\mu = -l_1 - l_3 + l_2 - \delta(r \leq i_1) - \delta(T \leq r + i_3)$ and $i \in \mathbb{Z}$. Then

$$\begin{aligned}
& \text{Res}_x \left((1+x)^{\text{wt } a-1+l_1+\delta(r \leq i_1)+r/T} x^{\mu-i} \right. \\
& \quad \times \sum_{j \in \mathbb{Z}} Y_M^{(r)}(a, b; \text{wt } a + \text{wt } b + m - n - 2 - j, j)(w) x^{-j-1} \Big) \\
&= \sum_{k=0}^{\infty} \binom{\text{wt } a - 1 + l_1 + \delta(r \leq i_1) + r/T}{k} \\
& \quad \times Y_M^{(r)}(a, b; \text{wt } a + \text{wt } b + m - n - 2 - \mu + i - k, \mu - i + k)(w) \\
&= \sum_{k=0}^{\infty} \binom{\text{wt } a - 1 + l_1 + \delta(r \leq i_1) + r/T}{k} \\
& \quad \times \text{Res}_{x_2} \text{Res}_{x_1-x_2} \left(x_2^{\text{wt } a + \text{wt } b + m - n - 2 - \mu + i - k} (x_1 - x_2)^{\mu - i + k} \right. \\
& \quad \times Y_M^{(r)}(a, b | x_2, x_1 - x_2)(w) \Big) \\
&= \text{Res}_{x_2} \text{Res}_{x_1-x_2} \left(x_1^{\text{wt } a-1+l_1+\delta(r \leq i_1)+r/T} x_2^{\text{wt } b-1+l_1+\delta(r^\vee \leq i_1)+r^\vee/T-l_2+i-1} \right. \\
& \quad \times (x_1 - x_2)^{-l_1-l_3+l_2-\delta(r \leq i_1)-\delta(T \leq r+i_3)-i} Y_M^{(r)}(a, b | x_2, x_1 - x_2)(w) \Big),
\end{aligned}$$

where we used (3.12) in the last step and $r^\vee$ is defined in (3.8). Thus, (5.5) becomes

$$\begin{aligned}
& o_{n,m}(a *_{n,p,m}^T b)w \\
&= \sum_{i=0}^{l_2} \binom{-l_1 - l_3 + l_2 - \delta(r \leq i_1) - \delta(T \leq r + i_3)}{i} \\
& \quad \times \text{Res}_{x_2} \text{Res}_{x_1-x_2} \left(x_1^{\text{wt } a-1+l_1+\delta(r \leq i_1)+r/T} x_2^{\text{wt } b-1+l_1+\delta(r^\vee \leq i_1)+r^\vee/T-l_2+i-1} \right. \\
& \quad \times (x_1 - x_2)^{-l_1-l_3+l_2-\delta(r \leq i_1)-\delta(T \leq r+i_3)-i} Y_M^{(r)}(a, b | x_2, x_1 - x_2)(w) \Big). \tag{5.6}
\end{aligned}$$

The rest of the proof is the same as that of [5, Lemma 5.1] by (2.16). $\square$

The following result is a direct consequence of Lemma 5.4.

Corollary 5.5. *If M is generated by one homogeneous element w as a (V, T) -module, then $M = \{a_i w \mid a \in V, i \in (1/T)\mathbb{Z}\}$.*

We define an $A_n^T(V)$ - $A_m^T(V)$ -bimodule structure on $\text{Hom}_{\mathbb{C}}(M(m), M(n))$ by

$$(afb)(w) = a(f(bw))$$

for $f \in \text{Hom}_{\mathbb{C}}(M(m), M(n))$, $a \in A_n^T(V)$, $b \in A_m^T(V)$ and $w \in M(m)$. For a (V, T) -module W and $m \in (1/T)\mathbb{N}$, define

$$\begin{aligned} \Omega_m(W) \\ = \{w \in W \mid a_{\text{wt } a-1+k} w = 0 \text{ for all homogeneous } a \in V \text{ and } k > m\}. \end{aligned}$$

Clearly, $\bigoplus_{i=0}^m M(i) \subset \Omega_m(M)$.

Lemmas 5.3 and 5.4 imply the following results.

Lemma 5.6. *For $u \in O_{n,m}^T(V)$, $o_{n,m}(u) = 0$ on $M(m)$. The linear map $o_{n,m} : V \rightarrow \text{Hom}_{\mathbb{C}}(M(m), M(n))$ induces an $A_n^T(V)$ - $A_m^T(V)$ -bimodule homomorphism from $A_{n,m}^T(V)$ to $\text{Hom}_{\mathbb{C}}(M(m), M(n))$.*

Proposition 5.7. *Let W be a (V, T) -module. Then $o : V \rightarrow \text{End}_{\mathbb{C}}(\Omega_m(W))$ induces a representation of $A_m^T(V)$. In particular, $M(m)$ is a left $A_m^T(V)$ -module.*

For a left $A_m^T(V)$ -module U , set

$$M(U) = \bigoplus_{n \in (1/T)\mathbb{N}} A_{n,m}^T(V) \otimes_{A_m^T(V)} U$$

and $M(U)(n) = A_{n,m}^T(V) \otimes_{A_m^T(V)} U$ for every $n \in (1/T)\mathbb{N}$. For homogeneous $a \in V$ and $i \in (1/T)\mathbb{Z}$, define an operator a_i from $M(U)(n)$ to $M(U)(n + \text{wt } a - i - 1)$ by

$$a_i(b \otimes u) = \begin{cases} (a *_{n+\text{wt } a-i-1, n, m}^T b) \otimes u & \text{if } n + \text{wt } a - i - 1 \geq 0, \\ 0 & \text{if } n + \text{wt } a - i - 1 < 0 \end{cases} \quad (5.7)$$

for $b \otimes u \in M(U)(n)$ with $b \in V$ and $u \in U$. This operation is well-defined (cf. [4, p.815]). We extend a_i for arbitrary $a \in V$ by linearity and set

$$Y_{M(U)}(a, x) = \sum_{i \in (1/T)\mathbb{Z}} a_i x^{-i-1} : M(U) \rightarrow M(U)((x^{1/T})).$$

We shall show $(M(U), Y_{M(U)})$ is a $(1/T)\mathbb{N}$ -graded (V, T) -module. For homogeneous $a, b \in V$, $s \in \mathbb{Z}$ with $0 \leq s \leq T-1$ and $i \in (1/T)\mathbb{Z}$, $j \in \mathbb{Z}$, define a linear map $Y_{M(U)}^{(s)}(a, b; i, j) : M(U)(n) \rightarrow M(U)(\text{wt } a + \text{wt } b - i - j - 2 + n)$ by

$$\begin{aligned} Y_{M(U)}^{(s)}(a, b; i, j)(c \otimes u) \\ = (\hat{E}_{\text{wt } a + \text{wt } b - i - j - 2 + n, n}^{(T; s)}(a, b, j) *_{\text{wt } a + \text{wt } b - i - j - 2 + n, n, m}^T c) \otimes u \end{aligned} \quad (5.8)$$

for $c \otimes u \in M(U)(n)$ with $c \in V$ and $u \in U$. This operation is also well-defined. We extend $Y_{M(U)}^{(s)}(a, b; i, j)$ for arbitrary $a, b \in V$ by linearity and set

$$Y_{M(U)}^{(s)}(a, b|x_2, x_0) = \sum_{i \in (1/T)\mathbb{Z}} \sum_{j \in \mathbb{Z}} Y_{M(U)}^{(s)}(a, b; i, j) x_2^{-i-1} x_0^{-j-1}.$$

It follows by (3.19) and (4.1) that $Y_{M(U)}^{(s)}(a, b|x_2, x_0)$ is a linear map from $M(U)$ to $M(U)((x_2^{1/T}))((x_0))$.

From now on, we simply write $M = M(U)$. By (4.5) we have

$$\begin{aligned} & \sum_{s=0}^{T-1} Y_M^{(s)}(a, b; i, j)(c \otimes u) \\ &= \sum_{s=0}^{T-1} (\hat{E}_{\text{wt } a + \text{wt } b - i - j - 2 + n, n}^{(T; s)}(a, b, j) *_{\text{wt } a + \text{wt } b - i - j - 2 + n, n, m}^T c) \otimes u \\ &= ((a_j b) *_{\text{wt } a + \text{wt } b - i - j - 2 + n, n, m}^T c) \otimes u \\ &= (a_j b)_i(c \otimes u) \end{aligned}$$

for homogeneous $a, b \in V$ and $c \otimes u \in M(n)$ with $c \in V$ and $u \in U$. Thus

$$\sum_{s=0}^{T-1} Y_M^{(s)}(a, b|x_2, x_0)(w) = Y_M(Y(a, x_0)b, x_2)w \quad (5.9)$$

for $w \in M$.

Lemma 5.8. (1) $a_i M(n) = 0$ for homogeneous $a \in V$ and $i > \text{wt } a - 1 + n$.

(2) $Y_M(\mathbf{1}, x) = \text{id}_M$.

Proof. Clearly, (1) holds. Let $a \otimes u \in M(n)$ with $a \in V$ and $c \in U$. By Lemma 3.6, we have

$$\begin{aligned} \mathbf{1}_i(a \otimes u) &= (\mathbf{1} *_{-i-1+n, n, m}^T a) \otimes u \\ &= \delta_{i, -1}(\mathbf{1}_{-1}a) \otimes u = \delta_{i, -1}a \otimes u \end{aligned}$$

for $i \in (1/T)\mathbb{Z}$. □

Lemma 5.9. *Let a, b be homogeneous elements of V , $i, j \in (1/T)\mathbb{Z}$ and r the integer uniquely determined by the conditions $0 \leq r \leq T-1$ and $r/T \equiv i \pmod{\mathbb{Z}}$. Then*

$$[a_i, b_j]w = \sum_{k=0}^{\infty} \binom{i}{k} Y_M^{(r)}(a, b; i+j-k, k)(w)$$

for $w \in M$. In particular,

$$(x_1 - x_2)^l [Y_M(a, x_1), Y_M(b, x_2)] = 0$$

for $l \in \mathbb{Z}_{\geq \text{wt } a + \text{wt } b - \Delta}$.

Proof. Let $c \otimes u \in M(n)$ with $c \in V$ and $u \in U$. By Lemma 4.3, we have

$$\begin{aligned} & a_i b_j (c \otimes u) - b_j a_i (c \otimes u) \\ &= (a *_{\text{wt } a + \text{wt } b - i - j - 2 + n, \text{wt } b - 1 - j + n, m}^T (b *_{\text{wt } b - 1 - j + n, n, m}^T c)) \otimes u \\ &\quad - (b *_{\text{wt } a + \text{wt } b - i - j - 2 + n, \text{wt } a - 1 - i + n, m}^T (a *_{\text{wt } a - 1 - i + n, n, m}^T c)) \otimes u \\ &= ((a *_{\text{wt } a + \text{wt } b - i - j - 2 + n, \text{wt } b - 1 - j + n, n}^T b) *_{\text{wt } a + \text{wt } b - i - j - 2 + n, n, m}^T c) \otimes u \\ &\quad - ((b *_{\text{wt } a + \text{wt } b - i - j - 2 + n, \text{wt } a - 1 - i + n, n}^T a) *_{\text{wt } a + \text{wt } b - i - j - 2 + n, n, m}^T c) \otimes u \\ &= (\text{Res}_x (1+x)^i \hat{X}_{\text{wt } a + \text{wt } b - i - j - 2 + n, n}^{(T; r)}(a, b; x) *_{\text{wt } a + \text{wt } b - i - j - 2 + n, n, m}^T c) \otimes u \\ &= \left(\sum_{k=0}^{\infty} \binom{i}{k} \hat{E}_{\text{wt } a + \text{wt } b - i - j - 2 + n, n}^{(T; r)}(a, b, k) *_{\text{wt } a + \text{wt } b - i - j - 2 + n, n, m}^T c \right) \otimes u \\ &= \sum_{k=0}^{\infty} \binom{i}{k} Y_M^{(r)}(a, b; i+j-k, k)(c \otimes u). \end{aligned}$$

The last formula follows from this and Remark 2.6 (cf. [13, Remark 3.1.13]). $\square$

We recall that $Y_M^r(a, x)$ denotes $\sum_{i \in r/T + \mathbb{Z}} a_i x^{-i-1}$ for $a \in V$ (cf. (2.10)).

Lemma 5.10. *Let $a, b \in V$ with a being homogeneous, $l, r \in \mathbb{N}$ with $0 \leq r \leq T-1$ and $n = l_3 + i_3/T \in (1/T)\mathbb{N}$ with $l_3, i_3 \in \mathbb{N}$ and $0 \leq i_3 \leq T-1$. Then*

$$\begin{aligned} & \text{Res}_{x_0} x_0^l (x_2 + x_0)^{\text{wt } a - 1 + l_3 + \delta(r \leq i_3) + r/T} Y_M^{(r)}(a, b | x_2, x_0)(w) \\ &= \text{Res}_{x_0} x_0^l (x_0 + x_2)^{\text{wt } a - 1 + l_3 + \delta(r \leq i_3) + r/T} Y_M^r(a, x_0 + x_2) Y_M(b, x_2) w \end{aligned}$$

for $w \in M(n)$.

Proof. Using Lemma 5.9, we obtain the formula by the same computation as in the proof of [5, Lemma 5.9]. $\square$

Lemma 5.11. *Let $a, b \in V$ with a being homogeneous, $r \in \mathbb{N}$ with $0 \leq r \leq T-1$ and $n = l_3 + i_3/T \in (1/T)\mathbb{N}$ with $l_3 \in \mathbb{N}$ and $0 \leq i_3 \leq T-1$. Then*

$$\begin{aligned} & \text{Res}_{x_0} x_0^{-l}(x_2 + x_0)^{\text{wt } a-1+l_3+\delta(r \leq i_3)+r/T} Y_M^{(r)}(a, b|x_2, x_0)(w) \\ &= \text{Res}_{x_0} x_0^{-l}(x_0 + x_2)^{\text{wt } a-1+l_3+\delta(r \leq i_3)+r/T} Y_M^r(a, x_0 + x_2) Y_M(b, x_2) w \end{aligned}$$

for $w \in M(n)$.

Proof. Let $c \otimes u \in M(n)$ with $c \in V$ and $u \in U$. We may assume b to be a homogeneous element of V . We shall show

$$\begin{aligned} & \text{Res}_{x_0} x_0^{-l}(x_2 + x_0)^{\text{wt } a+q} x_2^{\text{wt } b-q} Y_M^{(r)}(a, b|x_2, x_0)(c \otimes u) \\ &= \text{Res}_{x_0} x_0^{-l}(x_0 + x_2)^{\text{wt } a+q} x_2^{\text{wt } b-q} Y_M^r(a, x_0 + x_2) Y_M(b, x_2) c \otimes u, \end{aligned}$$

where $q = -1 + l_3 + \delta(r \leq i_3) + r/T$. We have

$$\begin{aligned} & \text{Res}_{x_0} x_0^{-l}(x_2 + x_0)^{\text{wt } a+q} x_2^{\text{wt } b-q} Y_M^{(r)}(a, b|x_2, x_0)(c \otimes u) \\ &= \sum_{j=0}^{\infty} \sum_{k \in (1/T)\mathbb{Z}} \binom{\text{wt } a-1+l_3+\delta(r \leq i_3)+r/T}{j} x_2^{-k-1+\text{wt } a+\text{wt } b-j} \\ & \quad \times Y_M^{(r)}(a, b; k, j-l)(c \otimes u) \\ &= \sum_{j=0}^{\infty} \sum_{k \in (1/T)\mathbb{N}} \binom{\text{wt } a-1+l_3+\delta(r \leq i_3)+r/T}{j} x_2^{-l+k-n+1} \\ & \quad \times Y_M^{(r)}(a, b; \text{wt } a + \text{wt } b - j + l - k + n - 2, j-l)(c \otimes u) \\ &= \sum_{j=0}^{\infty} \sum_{k \in (1/T)\mathbb{N}} \binom{\text{wt } a-1+l_3+\delta(r \leq i_3)+r/T}{j} x_2^{-l+k-n+1} \\ & \quad \times (\hat{E}_{k,n}^{(T;r)}(a, b, j-l) *_{k,n,m}^T c) \otimes u \\ &= \sum_{k \in (1/T)\mathbb{N}} x_2^{-l+k-n+1} (\text{Res}_x x^{-l}(1+x)^{\text{wt } a-1+l_3+\delta(r \leq i_3)+r/T} \\ & \quad \times \hat{X}_{k,n}^{(T;r)}(a, b; x)) *_{k,n,m}^T c \otimes u. \end{aligned} \tag{5.10}$$

On the other hand, applying the same computation as in the proof

of [5, Lemma 5.10] to

$$\begin{aligned} & \text{Res}_{x_0} x_0^{-l} (x_0 + x_2)^{\text{wt } a+q} x_2^{\text{wt } b-q} Y_M^r(a, x_0 + x_2) Y_M(b, x_2) (c \otimes u) \\ &= \sum_{s=0}^{T-1} \text{Res}_{x_0} x_0^{-l} (x_0 + x_2)^{\text{wt } a+q} x_2^{\text{wt } b-q} Y_M^r(a, x_0 + x_2) Y_M^s(b, x_2) (c \otimes u), \end{aligned}$$

we have

$$\begin{aligned} & \text{Res}_{x_0} x_0^{-l} (x_0 + x_2)^{\text{wt } a+q} x_2^{\text{wt } b-q} Y_M^r(a, x_0 + x_2) Y_M(b, x_2) (c \otimes u) \\ &= \sum_{s=0}^{T-1} \sum_{\substack{k \in (i_3-r-s)/T + \mathbb{Z} \\ 0 \leq k}} x_2^{-l+k-n+1} \\ & \quad \times \sum_{\substack{j \in (i_3-s)/T + \mathbb{Z} \\ 0 \leq j \leq k+l_3+\delta(r \leq i_3)+r/T-l}} \binom{-l}{-j+l_3+\delta(r \leq i_3)+r/T-l+k} \\ & \quad \times (-1)^{-j+l_3+\delta(r \leq i_3)+r/T-l+k} (a *_{k,j,m}^T (b *_{j,n,m}^T c)) \otimes u \\ &\equiv \sum_{s=0}^{T-1} \sum_{\substack{k \in (i_3-r-s)/T + \mathbb{Z} \\ 0 \leq k}} x_2^{-l+k-n+1} \\ & \quad \times \sum_{\substack{j \in (i_3-s)/T + \mathbb{Z} \\ 0 \leq j \leq k+l_3+\delta(r \leq i_3)+r/T-l}} \binom{-l}{-j+l_3+\delta(r \leq i_3)+r/T-l+k} \\ & \quad \times (-1)^{-j+l_3+\delta(r \leq i_3)+r/T-l+k} ((a *_{k,j,n}^T b) *_{k,n,m}^T c) \otimes u \\ & \quad (\text{mod } O_{n,m}^T(V)((x_2))). \end{aligned} \tag{5.11}$$

Moreover, for each $k = l_4 + (i_3 - r - s)/T \in (i_3 - r - s)/T + \mathbb{Z}$ with $k \geq 0$ and $l_4 \in \mathbb{Z}$, we have

$$\begin{aligned} & \sum_{\substack{j \in (i_3-s)/T + \mathbb{Z} \\ 0 \leq j \leq k+l_3+\delta(r \leq i_3)+r/T-l}} \binom{-l}{-j+l_3+\delta(r \leq i_3)+r/T-l+k} \\ & \quad \times (-1)^{-j+l_3+\delta(r \leq i_3)+r/T-l+k} a *_{k,j,n}^T b \\ &= \sum_{p=0}^{l_4+l_3+\delta(r \leq i_3)+\delta(s \leq i_3)-l-1} \binom{-l}{p} (-1)^p \sum_{i=0}^{l_4+l_3+\delta(r \leq i_3)+\delta(s \leq i_3)-l-1-p} \binom{-p-l}{i} \\ & \quad \times \text{Res}_x (1+x)^{\text{wt } a-1+l_3+\delta(r \leq i_3)+r/T} x^{-p-l-i} \hat{X}_{k,n}^{(T;r)}(a, b; x) \\ &= \text{Res}_x x^{-l} (1+x)^{\text{wt } a-1+l_3+\delta(r \leq i_3)+r/T} \hat{X}_{k,n}^{(T;r)}(a, b; x) \end{aligned} \tag{5.12}$$

by [4, Proposition 5.3]. By (5.10)–(5.12) the proof is complete. $\square$

By Lemmas 5.10 and 5.11, we have the following result.

Lemma 5.12. *Let $a, b \in V$ with a being homogeneous, $r \in \mathbb{N}$ with $0 \leq r \leq T - 1$ and $n = l_3 + i_3/T \in (1/T)\mathbb{N}$ with $l_3, i_3 \in \mathbb{N}$ and $0 \leq i_3 \leq T - 1$. Then*

$$\begin{aligned} & (x_2 + x_0)^{\text{wt } a - 1 + l_3 + \delta(r \leq i_3) + r/T} Y_M^{(r)}(a, b | x_2, x_0) \\ &= (x_0 + x_2)^{\text{wt } a - 1 + l_3 + \delta(r \leq i_3) + r/T} Y_M^r(a, x_0 + x_2) Y_M(b, x_2) \end{aligned}$$

on $M(n)$.

By (5.9) and Lemmas 2.4, 5.8, 5.9 and 5.12, the same argument as in the proof of [4, Theorem 4.13] shows the following theorem.

Theorem 5.13. *Let U be a left $A_m^T(V)$ -module. Then $M(U) = \bigoplus_{n \in (1/T)\mathbb{N}} A_{n,m}^T(V) \otimes_{A_m^T(V)} U$ is a $(1/T)\mathbb{N}$ -graded (V, T) -module with $M(U)(n) = A_{n,m}^T(V) \otimes_{A_m^T(V)} U$ and the following universal property: for a (V, T) -module W and an $A_m^T(V)$ -homomorphism $\sigma : U \rightarrow \Omega_m(W)$, there is a unique homomorphism $\bar{\sigma} : M(U) \rightarrow W$ of (V, T) -modules that extends σ . Moreover, if U cannot factor through $A_{m-1/T}^T(V)$, then $M(U)(0) \neq 0$.*

The following result immediately follows from Theorem 5.13 (cf. [7, Theorem 4.9]).

Corollary 5.14. *For every $m \in (1/T)\mathbb{N}$, there is a bijection between the set of isomorphism classes of simple left $A_m^T(V)$ -modules which cannot factor through $A_{m-1/T}^T(V)$ and that of simple $(1/T)\mathbb{N}$ -graded (V, T) -modules.*

6 Appendix

6.1 The determinant of a matrix

In this subsection we shall show that the matrix Γ in (3.4) is non-singular. Let b, t be positive integers and $x_0, \dots, x_{t-1}$ indeterminates. We denote by E_n the $n \times n$ identity matrix. Define $\alpha_i^k(x_s) = \sum_{j=1}^k \binom{x_s}{i+j} \binom{-x_s}{k-j} \in \mathbb{C}[x_s]$ for $0 \leq s \leq t-1, 1 \leq k \leq b$ and $i \in \mathbb{Z}$. Note that

$$\deg \alpha_i^k(x_s) = i + k \tag{6.1}$$

for $i \in \mathbb{N}$. Define t $bt \times b$ -matrices $A_s, s = 0, \dots, t-1$ by

$$A_s = \begin{pmatrix} \alpha_{(t-1)b-1}^1(x_s) & \alpha_{(t-1)b-1}^2(x_s) & \cdots & \alpha_{(t-1)b-1}^b(x_s) \\ \alpha_{(t-1)b-2}^1(x_s) & \alpha_{(t-1)b-2}^2(x_s) & \cdots & \alpha_{(t-1)b-2}^b(x_s) \\ \vdots & \vdots & & \vdots \\ \alpha_{-b}^1(x_s) & \alpha_{-b}^2(x_s) & \cdots & \alpha_{-b}^b(x_s) \end{pmatrix} \quad (6.2)$$

and set $A = (A_0 \cdots A_{t-1})$. The following result implies Γ is non-singular.

Proposition 6.1.

$$\det A = \prod_{0 \leq i < j \leq t-1} \prod_{k=-b+1}^{b-1} \left(\frac{x_i - x_j + k}{b(j-i) + k} \right)^{b-|k|}.$$

Proof. Since

$$\begin{aligned} & (\alpha_i^1(x_s), \dots, \alpha_i^b(x_s)) \\ &= \left(\binom{x_s}{i+1}, \binom{x_s}{i+2}, \dots, \binom{x_s}{i+b} \right) \begin{pmatrix} 1 & \binom{-x_s}{1} & \binom{-x_s}{2} & \cdots & \binom{-x_s}{b} \\ 0 & 1 & \binom{-x_s}{1} & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \binom{-x_s}{2} \\ \vdots & & \ddots & \ddots & \binom{-x_s}{1} \\ 0 & \cdots & \cdots & 0 & 1 \end{pmatrix}, \end{aligned}$$

the determinant of A is equal to that of $B = (B_0 \cdots B_{t-1})$, where

$$B_s = \begin{pmatrix} \binom{x_s}{(t-1)b} & \binom{x_s}{(t-1)b+1} & \cdots & \binom{x_s}{tb-1} \\ \binom{x_s}{(t-1)b-1} & \binom{x_s}{(t-1)b} & \cdots & \binom{x_s}{tb-2} \\ \vdots & \vdots & & \vdots \\ \binom{x_s}{-b+1} & \binom{x_s}{-b+2} & \cdots & \binom{x_s}{0} \end{pmatrix}, \quad s = 0, \dots, t-1.$$

The same argument as in the proof of [15, Proposition 9] shows that $(x_i - x_j + k)^{b-|k|}$ is a factor of $\det B$ for each $0 \leq i < j \leq t-1$ and $-b+1 \leq k \leq b-1$. Thus, there is $c \in \mathbb{C}[x_0, \dots, x_{t-1}]$ such that

$$\det B = c \prod_{0 \leq i < j \leq t-1} \prod_{k=-b+1}^{b-1} (x_i - x_j + k)^{b-|k|}.$$

Since $\alpha_i^k(x_s) = \delta_{i+k,0}$ for $i < 0$, we have $A_s = \begin{pmatrix} A'_s \\ E_b \end{pmatrix}$, $s = 0, \dots, T-1$, where

$$A'_s = \begin{pmatrix} \alpha_{(t-1)b-1}^1(x_s) & \alpha_{(t-1)b-1}^2(x_s) & \cdots & \alpha_{(t-1)b-1}^b(x_s) \\ \alpha_{(t-1)b-2}^1(x_s) & \alpha_{(t-1)b-2}^2(x_s) & \cdots & \alpha_{(t-1)b-2}^b(x_s) \\ \vdots & \vdots & & \vdots \\ \alpha_0^1(x_s) & \alpha_0^2(x_s) & \cdots & \alpha_0^b(x_s) \end{pmatrix}.$$

It follows by

$$\begin{pmatrix} E_{(t-1)b} & -A'_0 \\ O & E_b \end{pmatrix} A = \begin{pmatrix} O & A'_1 - A'_0 & \cdots & A'_{t-1} - A'_0 \\ E_b & E_b & \cdots & E_b \end{pmatrix} \quad (6.3)$$

that $\det A = (-1)^{(t-1)b^2} \det (A'_1 - A'_0 \cdots A'_{t-1} - A'_0)$. Thus, the degree of $\det A \in \mathbb{C}[x_0, \dots, x_{t-1}]$ is at most $\binom{t}{2}b^2$ by (6.1). Since the degree of $\prod_{0 \leq i < j \leq t-1} \prod_{k=-b+1}^{b-1} (x_i - x_j + k)^{b-|k|}$ is equal to $\binom{t}{2}b^2$, we have $c \in \mathbb{C}$.

Substituting $((t-1)b, (t-2)b, \dots, 0)$ for $(x_0, x_1, \dots, x_{t-1})$, we obtain

$$1 = c \prod_{0 \leq i < j \leq t-1} \prod_{k=-b+1}^{b-1} (b(j-i) + k)^{b-|k|}.$$

The proof is complete. $\square$

6.2 Some improvements of results on $A_{G,n}(V)$

The purpose of this subsection is to improve Theorems 1 and 2 in [15]. Let $V = \oplus_{j=\Delta}^{\infty} V_j$ be a vertex operator algebra and G an automorphism group of V of finite order t . For $g \in G$ and $n \in (1/t)\mathbb{N}$, $O_{g,n}(V)$ is the subspace of V defined in [8].

In [15], under the condition that $\Delta = 0$, we constructed an associative algebra $A_{G,n}(V)$ for each $n \in (1/t)\mathbb{Z}$ in Theorem 1 and got a duality theorem of Schur-Weyl type in Theorem 2 by using $A_{G,n}(V)$. The condition that $\Delta = 0$ was used in order to show the non-singularity of a matrix in [15, Lemma 3].

We shall show [15, Theorems 1 and 2] without assuming $\Delta = 0$. To do this, it is sufficient to show the following lemma, which is an improvement of [15, Lemma 3], by using $\hat{E}_{n,m}^{(t;s)}(a, b, i)$ defined in (4.1). We note that the existence of $\hat{E}_{n,m}^{(t;s)}(a, b, i)$ follows from Lemma 3.2 and Proposition 6.1. We use the notation in Remark 2.7.

Lemma 6.2. *For $a, b \in V = \oplus_{j=\Delta}^{\infty} V_j$, $0 \leq r \leq t-1$, $p \in \mathbb{Z}$, $n \in (1/t)\mathbb{N}$ and $g \in G$, we have*

$$\hat{E}_{n,n}^{(t;r)}(a, b, p) \equiv a_p^{(g,r)} b \pmod{O_{g,n}(V)}$$

Proof. We may assume a, b to be homogeneous. We write $n = l + i/t$ with $l, i \in \mathbb{N}$ and $0 \leq i \leq t-1$. Let $q = -2l - 3$ and $Q_s = \text{wt } a - 1 + l + \delta(s \leq i) + s/t$, $s = 0, \dots, t-1$. We use the notation in Section 3. It follows from (3.1) that the image of the subspace $O(\text{wt } a + \text{wt } b - 1 - \Delta, Q_s, q; z)$ of $\mathbb{C}[z, z^{-1}]$ under the map $\mathbb{C}[z, z^{-1}] \ni f \mapsto f|_{z^j=a_j^{(g,s)}b} \in V$ is contained in $O_{g,n}(V)$ for $s = 0, \dots, t-1$. By (3.18), we have

$$\begin{aligned} \hat{E}_{n,n}^{(t;r)}(a, b, p) &= E_{n,n}^{(t;r)}(\text{wt } a, \text{wt } b, p; z)|_{z^j=a_j b} \\ &= \sum_{s \neq r} E_{n,n}^{(t;r)}(\text{wt } a, \text{wt } b, p; z)|_{z^j=a_j^{(g,s)}b} \\ &\quad + E_{n,n}^{(t;r)}(\text{wt } a, \text{wt } b, p; z)|_{z^j=a_j^{(g,r)}b} \\ &\equiv a_p^{(g,r)} b \pmod{O_{g,n}(V)}. \end{aligned}$$

□

References

- [1] R. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster, *Proc. Nat. Acad. Sci. U.S.A.* **83** (1986), 3068–3071.
- [2] L. Dixon, J.A. Harvey, C. Vafa and E. Witten, String on orbifolds, *Nucl. Phys. B* **261** (1985) 620–678; String on orbifolds II, *Nucl. Phys. B* **274** (1986) 285–314.
- [3] R. Dijkgraaf, C. Vafa, E. Verlinde, and H. Verlinde, The operator algebra of orbifold models, *Comm. Math. Phys.* **123** (1989), 485–526.
- [4] C. Dong and C. Jiang, Bimodules associated to vertex operator algebras, *Math. Z.* **259** (2008), 799–826.
- [5] C. Dong and C. Jiang, Bimodules and g -rationality of vertex operator algebras, *Trans. Amer. Math. Soc.* **360** (2008), 4235–4262.
- [6] C. Dong, H.S. Li and G. Mason, Twisted representations of vertex operator algebras, *Math. Ann.* **310** (1998), 571–600.

- [7] C. Dong, H.S. Li and G. Mason, Vertex operator algebras and associative algebras, *J. Algebra* **206** (1998), 67–96.
- [8] C. Dong, H.S. Li and G. Mason, Twisted representations of vertex operator algebras and associative algebras, *Internat. Math. Res. Notices* (1998), no. 8, 389–397.
- [9] C. Dong and G. Mason, On quantum Galois theory. *Duke Math. J.* **86** (1997), 305–321.
- [10] I. B. Frenkel, J. Lepowsky and A. Meurman, *Vertex Operator Algebras and the Monster*, Pure and Applied Math., Vol. **134**, Academic Press, 1988.
- [11] A. Hanaki, M. Miyamoto and D. Tambara, Quantum Galois theory for finite groups. *Duke Math. J.* **97** (1999), 541–544.
- [12] J. Lepowsky, Calculus of twisted vertex operators, *Proc. Nat. Acad. Sci. U.S.A.* **82** (1985), 8295–8299.
- [13] J. Lepowsky and H. S. Li, *Introduction to Vertex Operator Algebras and their Representations*, Progress in Mathematics, **227**, Birkhauser Boston, Inc., Boston, MA, 2004.
- [14] H. S. Li, Local systems of twisted vertex operators, vertex operator superalgebras and twisted modules, *Contemp. Math.* **193** (1996), 203–236.
- [15] M. Miyamoto and K. Tanabe, Uniform product of $A_{g,n}(V)$ for an orbifold model V and G -twisted Zhu algebra. *J. Algebra* **274** (2004), 80–96.
- [16] Y. Zhu, Modular invariance of characters of vertex operator algebras, *J. Amer. Math. Soc.* **9** (1996), 237–302.